



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 39/395, 48/00, C12P 19/34, C12Q 1/68, G01N 33/53, 33/574, 33/546, 33/567		A1	(11) International Publication Number: WO 00/23111
			(43) International Publication Date: 27 April 2000 (27.04.00)
(21) International Application Number: PCT/US99/24331		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 19 October 1999 (19.10.99)			
(30) Priority Data: 60/104,737 19 October 1998 (19.10.98) US		Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant (for all designated States except US): DIADEXUS LLC [US/US]; 3303 Octavius Drive, Santa Clara, CA 95054 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): SALCEDA, Susana [AR/US]; 4118 Crescendo Avenue, San Jose, CA 95136 (US). RECIPON, Herve [FR/US]; 85 Fortuna Avenue, San Francisco, CA 94115 (US). CAFFERKEY, Robert [IE/US]; Apartment #218, 350 Elan Village Lane, San Jose, CA 95134 (US).			
(74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).			
(54) Title: METHOD OF DIAGNOSING, MONITORING, STAGING, IMAGING AND TREATING PROSTATE CANCER			
(57) Abstract The present invention provides new methods for detecting, diagnosing, monitoring, staging, prognosticating, imaging and treating prostate cancer.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LJ	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

**METHOD OF DIAGNOSING,
MONITORING, STAGING, IMAGING AND TREATING PROSTATE CANCER**

FIELD OF THE INVENTION

This invention relates, in part, to newly developed
5 assays for detecting, diagnosing, monitoring, staging,
prognosticating, imaging and treating cancers, particularly
prostate cancer.

BACKGROUND OF THE INVENTION

Cancer of the prostate is the most prevalent malignancy
10 in adult males, excluding skin cancer, and is an increasingly
prevalent health problem in the United States. In 1996, it
was estimated that 41,400 deaths would result from this
disease in the United States alone, indicating that prostate
cancer is second only to lung cancer as the most common cause
15 of death in the same population. If diagnosed and treated
early, when the cancer is still confined to the prostate, the
chances of cure is significantly higher.

Treatment decisions for an individual are linked to the
stage of prostate cancer present in that individual. A common
20 classification of the spread of prostate cancer was developed
by the American Urological Association (AUA). The AUA system
divides prostate tumors into four stages, A to D. Stage A,
microscopic cancer within prostate, is further subdivided into
stages A1 and A2. Sub-stage A1 is a well-differentiated
25 cancer confined to one site within the prostate. Treatment
is generally observation, radical prostatectomy, or radiation.
Sub-stage A2 is a moderately to poorly differentiated cancer
at multiple sites within the prostate. Treatment is radical
prostatectomy or radiation. Stage B, palpable lump within the
30 prostate, is also further subdivided into sub-stages B1 and
B2. In sub-stage B1, the cancer forms a small nodule in one

- 2 -

lobe of the prostate. In sub-stage B2, the cancer forms large or multiple nodules, or occurs in both lobes of the prostate. Treatment for sub-stages B1 and B2 is either radical prostatectomy or radiation. Stage C is a large cancer mass
5 involving most or all of the prostate and is also further subdivided into two sub-stages. In sub-stage C1, the cancer forms a continuous mass that may have extended beyond the prostate. In sub-stage C2, the cancer forms a continuous mass that invades the surrounding tissue. Treatment for both these
10 sub-stages is radiation with or without drugs to address the cancer. The fourth stage, Stage D is metastatic cancer and is also subdivided into two sub-stages. In sub-stage D1, the cancer appears in the lymph nodes of the pelvis. In sub-stage D2, the cancer involves tissues beyond lymph nodes. Treatment
15 for both of these sub-stages is systemic drugs to address the cancer as well as pain.

However, current prostate cancer staging methods are limited. As many as 50% of prostate cancers initially staged as A2, B, or C are actually stage D, metastatic. Discovery
20 of metastasis is significant because patients with metastatic cancers have a poorer prognosis and require significantly different therapy than those with localized cancers. The five year survival rates for patients with localized and metastatic prostate cancers are 93% and 29%, respectively.

25 Accordingly, there is a great need for more sensitive and accurate methods for the staging of a cancer in a human to determine whether or not such cancer has metastasized and for monitoring the progress of a cancer in a human which has not metastasized for the onset of metastasis.

30 It has now been found that a number of proteins in the public domain are useful as diagnostic markers for prostate cancer. These diagnostic markers are referred to herein as cancer specific genes or CSGs and include, but are not limited to: Pro109 which is a human zinc- α 2-glycoprotein (Freje et
35 al. Genomics 1993 18(3):575-587); Pro112 which is a human

- 3 -

cysteine-rich protein with a zinc-finger motif (Liebhaber et al. Nucleic Acid Research 1990 18(13):3871-3879; WO9514772 and WO9845436); Prol11 which is a prostate-specific transglutaminase (Dubbink et al. Genomics 1998 51(3):434-444);
5 Prol15 which is a novel serine protease with transmembrane, LDLR, and SRCR domains and maps to 21q22.3 (Paoloni-Giacobino et al. Genomics 1997 44(3):309-320; WO9837418 and WO987093); Prol10 which is a human breast carcinoma fatty acid synthase (U.S. Patent 5,665,874 and WO9403599); Prol13 which is a
10 homeobox gene, HOXB13 (Steinicki et al. J. Invest. Dermatol. 1998 111:57-63); Prol14 which is a human tetraspan NET-1 (WO9839446); and Prol18 which is a human JM27 protein (WO9845435). ESTs for these CSGs are set forth in SEQ ID NO: 1, 3, 5, 7, 9, 11, 13 and 15 while the full length contigs for
15 these CSGs are set forth in SEQ ID NO:2, 4, 6, 8, 10, 12, 14 and 16, respectively. Additional CSGs for use in the present invention are depicted herein in SEQ ID NO: 17, 18, 19 and 20.

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging, prognosticating,
20 imaging and treating prostate cancer via the cancer specific genes referred to herein as CSGs. For purposes of the present invention, CSG refers, among other things, to native protein expressed by the gene comprising a polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15,
25 16, 17, 18, 19 or 20. By "CSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1-20, but which still encode the same protein. In the alternative, what is meant by CSG as used herein, means the
30 native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, levels of the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or
35 20, or levels of a polynucleotide which is capable of

- 4 -

hybridizing under stringent conditions to the antisense sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein a change in levels of CSG in the patient versus the normal human control is associated with prostate cancer.

Further provided is a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which is not known to have metastasized by identifying a human patient suspected of having prostate cancer that has metastasized; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control, wherein an increase in CSG levels in the patient versus the normal human control is associated with prostate cancer which has metastasized.

- 5 -

Also provided by the invention is a method of staging prostate cancer in a human which has such cancer by identifying a human patient having such cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient
5 for CSG; comparing CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer
10 which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

Further provided is a method of monitoring prostate cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient
15 having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of
20 a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided is a method of monitoring the change in stage of prostate cancer in a human having such cancer by
25 looking at levels of CSG in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of
30 CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is
35 regressing or in remission.

- 6 -

Further provided are methods of designing new therapeutic agents targeted to a CSG for use in imaging and treating prostate cancer. For example, in one embodiment, therapeutic agents such as antibodies targeted against CSG or fragments of such antibodies can be used to detect or image localization of CSG in a patient for the purpose of detecting or diagnosing a disease or condition. Such antibodies can be polyclonal, monoclonal, or omniconal or prepared by molecular biology techniques. The term "antibody", as used herein and throughout the instant specification is also meant to include aptamers and single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art. Antibodies can be labeled with a variety of detectable labels including, but not limited to, radioisotopes and paramagnetic metals. Therapeutics agents such as antibodies or fragments thereof can also be used in the treatment of diseases characterized by expression of CSG. In these applications, the antibody can be used without or with derivatization to a cytotoxic agent such as a radioisotope, enzyme, toxin, drug or a prodrug.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging and prognosticating cancers

- 7 -

by comparing levels of CSG in a human patient with those of CSG in a normal human control. For purposes of the present invention, what is meant by CSG levels is, among other things, native protein expressed by the gene comprising a polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. By "CSG" it is also meant herein polynucleotides which, due to degeneracy in genetic coding, comprise variations in nucleotide sequence as compared to SEQ ID NO: 1-20, but which still encode the same protein. The native protein being detected, may be whole, a breakdown product, a complex of molecules or chemically modified. In the alternative, what is meant by CSG as used herein, means the native mRNA encoded by the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, levels of the gene comprising the polynucleotide sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20, or levels of a polynucleotide which is capable of hybridizing under stringent conditions to the antisense sequence of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20. Such levels are preferably determined in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for diagnosing overexpression of CSG protein compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of prostate cancer.

All the methods of the present invention may optionally include determining the levels of other cancer markers as well as CSG. Other cancer markers, in addition to CSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

- 8 -

Diagnostic Assays

The present invention provides methods for diagnosing the presence of prostate cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of CSG in the patient versus the normal human control is associated with the presence of prostate cancer.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic prostate cancer in a patient having prostate cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a human cancer patient suspected of having prostate cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art.

In the present invention, determining the presence of CSG levels in cells, tissues or bodily fluid, is particularly useful for discriminating between prostate cancer which has not metastasized and prostate cancer which has metastasized. Existing techniques have difficulty discriminating between prostate cancer which has metastasized and prostate cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, the cancer marker levels measured in such cells, tissues or bodily fluid is CSG, and are compared with levels of CSG in preferably the same cells, tissue or bodily fluid type of a normal human control. That

- 9 -

is, if the cancer marker being observed is just CSG in serum, this level is preferably compared with the level of CSG in serum of a normal human control. An increase in the CSG in the patient versus the normal human control is associated with prostate cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues or bodily fluid of a normal patient.

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may preferably also include samples from a human patient that is determined by reliable methods to have prostate cancer which has not metastasized.

20 **Staging**

The invention also provides a method of staging prostate cancer in a human patient. The method comprises identifying a human patient having such cancer and analyzing cells, tissues or bodily fluid from such human patient for CSG. The CSG levels determined in the patient are then compared with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG (but still increased over true normal levels) is associated with a cancer which is regressing or in remission.

Monitoring

Further provided is a method of monitoring prostate cancer in a human patient having such cancer for the onset of

- 10 -

metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; and comparing the CSG levels
5 determined in the human patient with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus the normal human control is associated with a cancer which has metastasized. In this method, normal
10 human control samples may also include prior patient samples.

Further provided by this invention is a method of monitoring the change in stage of prostate cancer in a human patient having such cancer. The method comprises identifying
15 a human patient having such cancer; periodically analyzing cells, tissues or bodily fluid from such human patient for CSG; and comparing the CSG levels determined in the human patient with levels of CSG in preferably the same cells, tissues or bodily fluid type of a normal human control, wherein an increase in CSG levels in the human patient versus
20 the normal human control is associated with a cancer which is progressing in stage and a decrease in the levels of CSG is associated with a cancer which is regressing in stage or in remission. In this method, normal human control samples may also include prior patient samples.

25 Monitoring a patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

Assay Techniques

30 Assay techniques that can be used to determine levels of gene expression (including protein levels), such as CSG of the present invention, in a sample derived from a patient are well known to those of skill in the art. Such assay methods include, without limitation, radioimmunoassays, reverse
35 transcriptase PCR (RT-PCR) assays, immunohistochemistry

- 11 -

assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses, ELISA assays and proteomic approaches: two-dimensional gel electrophoresis (2D electrophoresis) and non-gel based approaches such as mass spectrometry or protein interaction profiling. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids.

An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to CSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to CSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to CSG is incubated on a solid support, e.g. a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time CSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to CSG and linked to a detectable reagent such as horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to CSG. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to CSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of CSG protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

- 12 -

A competition assay can also be employed wherein antibodies specific to CSG are attached to a solid support and labeled CSG and a sample derived from the host are passed over the solid support. The amount of label detected which is
5 attached to the solid support can be correlated to a quantity of CSG in the sample.

Nucleic acid methods can also be used to detect CSG mRNA as a marker for prostate cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain
10 reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population
15 in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the
20 presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on
25 a solid support (i.e. gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the CSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or
30 plastic. At least a portion of the DNA encoding the CSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest. Hybridization between the substrate bound DNA and the analyte
35 can be detected and quantitated by several means including but

- 13 -

not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the
5 analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

Of the proteomic approaches, 2D electrophoresis is a
10 technique well known to those in the art. Isolation of individual proteins from a sample such as serum is accomplished using sequential separation of proteins by different characteristics usually on polyacrylamide gels. First, proteins are separated by size using an electric
15 current. The current acts uniformly on all proteins, so smaller proteins move farther on the gel than larger proteins. The second dimension applies a current perpendicular to the first and separates proteins not on the basis of size but on the specific electric charge carried by each protein. Since
20 no two proteins with different sequences are identical on the basis of both size and charge, the result of a 2D separation is a square gel in which each protein occupies a unique spot. Analysis of the spots with chemical or antibody probes, or subsequent protein microsequencing can reveal the relative
25 abundance of a given protein and the identity of the proteins in the sample.

The above tests can be carried out on samples derived from a variety of cells, bodily fluids and/or tissue extracts such as homogenates or solubilized tissue obtained from a
30 patient. Tissue extracts are obtained routinely from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva or any other bodily secretion or derivative thereof. By blood it is meant to include whole blood, plasma, serum or any derivative of
35 blood.

- 14 -

In Vivo Targeting of CSGs

Identification of these CSGs is also useful in the rational design of new therapeutics for imaging and treating cancers, and in particular prostate cancer. For example, in one embodiment, antibodies which specifically bind to CSG can be raised and used *in vivo* in patients suspected of suffering from prostate cancer. Antibodies which specifically bind a CSG can be injected into a patient suspected of having prostate cancer for diagnostic and/or therapeutic purposes.

The preparation and use of antibodies for *in vivo* diagnosis is well known in the art. For example, antibody-chelators labeled with Indium-111 have been described for use in the radioimmunoscinotographic imaging of carcinoembryonic antigen expressing tumors (Sumerdon et al. Nucl. Med. Biol. 1990 17:247-254). In particular, these antibody-chelators have been used in detecting tumors in patients suspected of having recurrent colorectal cancer (Griffin et al. J. Clin. Onc. 1991 9:631-640). Antibodies with paramagnetic ions as labels for use in magnetic resonance imaging have also been described (Lauffer, R.B. Magnetic Resonance in Medicine 1991 22:339-342). Antibodies directed against CSG can be used in a similar manner. Labeled antibodies which specifically bind CSG can be injected into patients suspected of having prostate cancer for the purpose of diagnosing or staging of the disease status of the patient. The label used will be selected in accordance with the imaging modality to be used. For example, radioactive labels such as Indium-111, Technetium-99m or Iodine-131 can be used for planar scans or single photon emission computed tomography (SPECT). Positron emitting labels such as Fluorine-19 can be used in positron emission tomography. Paramagnetic ions such as Gadlinium (III) or Manganese (II) can be used in magnetic resonance imaging (MRI). Localization of the label permits determination of the spread of the cancer. The amount of label within an organ or

- 15 -

tissue also allows determination of the presence or absence of cancer in that organ or tissue.

For patients diagnosed with prostate cancer, injection of an antibody which specifically binds CSG can also have a therapeutic benefit. The antibody may exert its therapeutic effect alone. Alternatively, the antibody can be conjugated to a cytotoxic agent such as a drug, toxin or radionuclide to enhance its therapeutic effect. Drug monoclonal antibodies have been described in the art for example by Garnett and Baldwin, Cancer Research 1986 46:2407-2412. The use of toxins conjugated to monoclonal antibodies for the therapy of various cancers has also been described by Pastan et al. Cell 1986 47:641-648. Yttrium-90 labeled monoclonal antibodies have been described for maximization of dose delivered to the tumor while limiting toxicity to normal tissues (Goodwin and Meares Cancer Supplement 1997 80:2675-2680). Other cytotoxic radionuclides including, but not limited to Copper-67, Iodine-131 and Rhenium-186 can also be used for labeling of antibodies against CSG.

Antibodies which can be used in these *in vivo* methods include polyclonal, monoclonal and omniclonal antibodies and antibodies prepared via molecular biology techniques. Antibody fragments and aptamers and single-stranded oligonucleotides such as those derived from an *in vitro* evolution protocol referred to as SELEX and well known to those skilled in the art can also be used.

Small molecules predicted via computer imaging to specifically bind to regions of CSGs can also be designed and synthesized and tested for use in the imaging and treatment of prostate cancer. Further, libraries of molecules can be screened for potential anticancer agents by assessing the ability of the molecule to bind to CSGs identified herein. Molecules identified in the library as being capable of binding to CSG are key candidates for further evaluation for use in the treatment of prostate cancer.

- 16 -

EXAMPLES

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments.

5 These exemplifications, while illustrating certain aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

All examples outlined here were carried out using standard techniques, which are well known and routine to those of skill in the art; except where otherwise described in detail. Routine molecular biology techniques of the following example can be carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

10
15

Example 1: Identification of CSGs

Identification of CSGs were carried out by a systematic analysis of data in the LIFESEQ database available from Incyte Pharmaceuticals, Palo Alto, CA, using the data mining Cancer Leads Automatic Search Package (CLASP) developed by diaDexus LLC, Santa Clara, CA.

20

The CLASP performs the following steps: selection of highly expressed organ specific genes based on the abundance level of the corresponding EST in the targeted organ versus all the other organs; analysis of the expression level of each highly expressed organ specific genes in normal, tumor tissue, disease tissue and tissue libraries associated with tumor or disease; selection of the candidates demonstrating component ESTs were exclusively or more frequently found in tumor libraries. The CLASP allows the identification of highly expressed organ and cancer specific genes. A final manual in depth evaluation is then performed to finalize the CSGs selection.

25
30

- 17 -

Clones depicted in the following Table 1 are CSGs useful in diagnosing, monitoring, staging, imaging and treating prostate cancer.

Table 1: CSGs

5	Clone ID	Pro #	SEQ ID NO:
	3424528H1	Pro109	1,2
	578349H1	Pro112	3,4
	1794013H1	Pro111	5,6
	2189835H1	Pro115	7,8
10	3277219H1	Pro110	9,10
	1857415	Pro113	11,12
	1810463H1	Pro114	13,14
	zr65G11	Pro118	15,16
	2626135H1		17
15	zd46d08		18
	1712252H1		19
	784583H1		20

Example 2: Relative Quantitation of Gene Expression

20 Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye.

25 During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

Amplification of an endogenous control is used to
 30 standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), ATPase, or 18S ribosomal RNA (rRNA) is used as this endogenous

- 18 -

control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample were used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained
5 using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

The tissue distribution and the level of the target gene were evaluated for every sample in normal and cancer tissues. Total RNA was extracted from normal tissues, cancer tissues,
10 and from cancers and the corresponding matched adjacent tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was done using primers and Taqman probes specific to each target gene. The results were analyzed using the ABI PRISM 7700
15 Sequence Detector. The absolute numbers are relative levels of expression of the target gene in a particular tissue compared to the calibrator tissue.

Expression of Clone ID 3424528H1 (Pro109):

For the CSG Pro109, real-time quantitative PCR was
20 performed using the following primers:

Forward Primer:

5'- ATCAGAACAAAGAGGCTGTGTC - 3' (SEQ ID NO:21)

Reverse Primer:

5'- ATCTCTAAAGCCCCAACCTTC - 3' (SEQ ID NO:22)

25 The absolute numbers depicted in Table 2 are relative levels of expression of the CSG referred to as Pro109 in 12 normal different tissues. All the values are compared to normal stomach (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular
30 tissue from different individuals.

- 19 -

Table 2: Relative Levels of CSG Prol09 Expression in Pooled Samples

Tissue	NORMAL
Colon	0.02
Endometrium	0.01
Kidney	0.48
Liver	14.83
Ovary	0.08
Pancreas	4.38
Prostate	11.24
Small Intestine	0.42
Spleen	0
Stomach	1
Testis	0.62
Uterus	0.02

The relative levels of expression in Table 2 show that with the exception of liver (14.83), Prol09 mRNA expression is higher (11.24) in prostate compared with all other normal tissues analyzed. Pancreas, with a relative expression level of 4.38, is the only other tissue expressing considerable mRNA for Prol09.

The absolute numbers in Table 2 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 3.

The absolute numbers depicted in Table 3 are relative levels of expression of Prol09 in 28 pairs of matching samples and 4 unmatched samples. All the values are compared to normal stomach (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

- 20 -

Table 3: Relative Levels of CSG Pro109 Expression in Individual Samples

	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro34B	Prostate 1	5.98	6.06
5	Pro65XB	Prostate 2	16.68	3.85
	Pro69XB	Prostate 3	20.46	6.82
	Pro78XB	Prostate 4	1.39	1.4
	Pro101XB	Prostate 5	24.8	9.8
	Pro12B	Prostate 6	9.1	0.2
10	Pro13XB	Prostate 7	0.5	9.7
	Pro20XB	Prostate 8	13	12.5
	Pro23B	Prostate 9	16.8	3
	Ovr100050	Ovary 1	0.4	
	Ovr1028	Ovary 2	1.9	
15	Ovr18GA	Ovary 3		0.1
	Ovr206I	Ovary 4		0.1
	Mam12X	Mammary Gland 1	13.5	1.4
	Mam47XP	Mammary Gland 2	0.7	0.2
	Lng47XQ	Lung 1	2.36	0.03
20	Lng60XL	Lung 2	7.39	0.2
	Lng75XC	Lung 3	0.77	0.27
	StoAC44	Stomach 1	0.05	1.19
	StoAC93	Stomach 2	0.55	0.8
	StoAC99	Stomach 3	0.12	3.04
25	ColAS43	Colon 1	16.11	0.07
	ColAS45	Colon 2	0.11	0.08
	ColAS46	Colon 3	4.99	0.4
	Liv15XA	Liver 1	8.43	10.97
	Liv42X	Liver 2	1.57	20.82

- 21 -

Liv94XA	Liver 3	2.98	9.19
Pan77X	Pancreas 1	36	32
Pan82XP	Pancreas 2	0.09	7.09
Pan92X	Pancreas 3	0.7	0
Pan71XL	Pancreas 4	2.48	0.73
Pan10343	Pancreas 5	46	5.5

0 = Negative

In the analysis of matching samples, the higher levels of expression were in prostate, showing a high degree of tissue specificity for prostate tissue. Of all the samples different than prostate analyzed, only 4 cancer samples (the cancer sample mammary 1 with 13.5, colon 1 with 16.11, liver 1 with 8.43, and lung 2 with 7.39) showed an expression comparable to the mRNA expression in prostate. These results confirmed some degree of tissue specificity as obtained with the panel of normal pooled samples (Table 2).

Furthermore, the level of mRNA expression was compared in cancer samples and the isogenic normal adjacent tissue from the same individual. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 3 shows overexpression of Prol09 in 6 out of 9 primary prostate cancer tissues compared with their respective normal adjacents. Thus, overexpression in the cancer tissue was observed in 66.66% of the prostate matching samples tested (total of 9 prostate matching samples).

Altogether, the degree of tissue specificity, plus the mRNA overexpression in 66.66% of the primary prostate matching samples tested is indicative of Prol09 being a diagnostic marker for prostate cancer.

- 22 -

Expression of Clone ID 578349H1 (Pro112):

For the CSG Pro112, real-time quantitative PCR was performed using the following primers:

Forward Primer

5' - TGCCGAAGAGGTTTCAGTGC - 3' (SEQ ID NO:23)

Reverse Primer

5' - GCCACAGTGGTACTGTCCAGAT - 3' (SEQ ID NO:24)

The absolute numbers depicted in Table 4 are relative levels of expression of the CSG Pro112 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 4: Relative Levels of CSG Pro112 Expression in Pooled Samples

Tissue	NORMAL
Brain	2.9
Heart	0.1
Kidney	0.2
Liver	0.2
Lung	7.7
Mammary	4.2
Muscle	0.1
Prostate	5.5
Small Intestine	1.8
Testis	1
Thymus	1
Uterus	21

The relative levels of expression in Table 4 show that Pro112 mRNA expression is the 3rd most highly expressed gene (after uterus and mammary) in the pool of normal prostate tissue compared to a total of 12 tissues analyzed. The absolute numbers in Table 4 were obtained analyzing pools of samples of a particular tissue from different individuals. These results demonstrate that Pro112 mRNA expression is specific for prostate thus indicating Pro112 to be a diagnostic marker for prostate disease especially cancer.

- 23 -

Expression of Clone ID 1794013H1 (Prol11):

For the CSG Prol11, real-time quantitative PCR was performed using the following primers:

Forward Primer

5' - GCTGCAAGTTCTCCACATTGA - 3' (SEQ ID NO:25)

Reverse Primer

5' - CAGCCGCAGGTGAAACAC - 3' (SEQ ID NO:26)

The absolute numbers depicted in Table 5 are relative levels of expression of the CSG Prol11 in 12 normal different tissues. All the values are compared to normal testis (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 5: Relative Levels of CSG Prol11 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.04
Heart	0
Kidney	0
Liver	0
Lung	0.05
Mammary	0.14
Muscle	5166.6
Prostate	1483.72
Small Intestine	0.33
Testis	1
Thymus	0.49
Uterus	0.07

The relative levels of expression in Table 5 show that Prol11 mRNA expression is extraordinarily high in the pool of normal prostate (1483.72) compared to all the other tissues analyzed with the exception of muscle (5166.6). These results demonstrate that Prol11 mRNA expression shows specificity for prostate and muscle.

The absolute numbers in Table 5 were obtained analyzing pools of samples of a particular tissue from different

- 24 -

individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 6.

The absolute numbers depicted in Table 6 are relative levels of expression of Prol11 in 48 pairs of matching and 18 unmatched samples. All the values are compared to normal testis (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 6: Relative Levels of CSG Prol11 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Pro101XB	Prostate 1	8.3	21.8
Pro12B	Prostate 2	2336	133
Pro13XB	Prostate 3	3.4	23
Pro20XB	Prostate 4	21.6	121.5
Pro23B	Prostate 5	19.4	3.7
Pro34B	Prostate 6	15	39
Pro65XB	Prostate 7	8	867
Pro69XB	Prostate 8	56	94
Pro78XB	Prostate 9	24	1515
Pro84XB	Prostate 10	119	15.35
Pro90XB	Prostate 11	8.08	112.2
Pro91XB	Prostate 12	0.88	51.8
ProC215	Prostate 13	0.3	
ProC234	Prostate 14	0.35	
ProC280	Prostate 15	436.5	
Pro109XB	Prostate 16	3.43	265
Pro110	Prostate 17	18.2	8.73

- 25 -

	Pro125XB	Prostate 18	0.34	186
	Pro326	Prostate 19	1392	110
	Pro10R	Prostate 20 (prostatitis)	0.5	
	Pro20R	Prostate 21 (prostatitis)	24.1	
5	Pro258	Prostate 22 (BPH)	4610	
	Pro263C	Prostate 23 (BPH)	0	
	Pro267A	Prostate 24 (BPH)	1.46	
	Pro271A	Prostate 25 (BPH)	0	
	Pro460Z	Prostate 26 (BPH)	1.47	
10	ProC032	Prostate 27 (BPH)	14.4	
	Tst39X	Testis 1	0	0
	Bld32XK	Bladder 1	0.44	0.41
	Bld46XK	Bladder 2	0	0
	Bld66X	Bladder 3	0	0
15	BldTR14	Bladder 4	0	0
	Kid106XD	Kidney 1	0	0
	Kid107XD	Kidney 2	0	0
	Kid109XD	Kidney 3	0	0
	Pan10343	Pancreas 1	0	0
20	Pan71XL	Pancreas 2	0	0
	Pan77X	Pancreas 3	0	0
	Liv15XA	Liver 1	0	0
	Liv42X	Liver 2	0	0
	ClnAS43	Colon 1	0	0
25	ClnAS45	Colon 2	0	0
	ClnAS46	Colon 3	0	0
	ClnAS67	Colon 4	0	0
	ClnAC19	Colon 5	0	0
	ClnAS12	Colon 6	0	0

- 26 -

	SmI21XA	Small Intestine 1	0	0
	SmIH89	Small Intestine 2	0	0
	Lng47XQ	Lung 1	0.7	0
	Lng60XL	Lung 2	0	0
5	Lng75XC	Lung 3	0	0
	Lng90X	Lung 4	0	0
	Mam12X	Mammary Gland 1	0	1.4
	Mam59X	Mammary Gland 2	0.2	0
	MamA06X	Mammary Gland 3	0	0
10	MamS127	Mammary Gland 4	0	0
	Mam162X	Mammary Gland 5	0	0
	Mam42DN	Mammary Gland 6	0	0
	Ovr103X	Ovary 1	0.14	0
	Ovr10050	Ovary 2	0.2	
15	Ovr1028	Ovary 3	0	
	Ovr10400	Ovary 4	0.2	
	Ovr18GA	Ovary 5		0
	Ovr206I	Ovary 6		0
	Ovr20GA	Ovary 7		0.2
20	Ovr25GA	Ovary 8		0

0= Negative

In the analysis of matching samples, the higher levels of expression were in prostate showing a high degree of tissue specificity for prostate. These results confirm the tissue specificity results obtained with normal pooled samples (Table 5).

Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 6 shows overexpression of Prol11 in 5 out

- 27 -

of 16 primary prostate cancer samples compared with their respective normal adjacent (prostate samples 2, 5, 10, 17, and 19). Similar expression levels were observed in 3 unmatched prostate cancers (prostate samples 13, 14, 15), 2 prostatitis
 5 (prostate samples 20, 21), and 6 benign prostatic hyperplasia samples (prostate samples 22 through 27). Thus, there is overexpression in the cancer tissue of 31.25% of the prostate matching samples tested (total of 16 prostate matching samples).

10 Altogether, the high level of tissue specificity, plus the mRNA overexpression in 31.25% of the prostate matching samples tested are indicative of Prol11 being a diagnostic marker for prostate cancer.

Expression of Clone ID 2189835H1 (Prol15):

15 For the CSG Prol15, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- TGGCTTTGAACTCAGGGTCA - 3' (SEQ ID NO:27)

Reverse Primer

20 5'- CGGATGCACCTCGTAGACAG - 3' (SEQ ID NO:28)

The absolute numbers depicted in Table 7 are relative levels of expression of the CSG Prol15 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available
 25 pools, originated by pooling samples of a particular tissue from different individuals.

Table 7: Relative Levels of CSG Prol15 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.016
Heart	0.002
Kidney	8.08
Liver	2.20
Lung	112.99

- 28 -

5	Mammary	29.45
	Muscle	0.05
	Prostate	337.79
	Small Intestine	7.54
	Testis	1.48
	Thymus	1
	Uterus	1.4

The relative levels of expression in Table 7 show that Prol15 mRNA expression is higher (337.79) in prostate compared with all the other normal tissues analyzed. Lung, with a relative expression level of 112.99, and mammary (29.446) are the other tissues expressing moderate levels of mRNA for Prol15. These results establish Prol15 mRNA expression to be highly specific for prostate.

The absolute numbers in Table 7 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 8.

The absolute numbers depicted in Table 8 are relative levels of expression of Prol15 in 17 pairs of matching and 21 unmatched samples. All the values are compared to normal thymus (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 8: Relative Levels of CSG Prol15 Expression in Individual Samples

30	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro12B	Prostate 1	1475.9	190.3
	ProC234	Prostate 2	169.61	
	Pro109XB	Prostate 3		639.53
	Pro101XB	Prostate 4	1985.2	2882.9

- 29 -

	Pro13XB	Prostate 5	34.9	13.9
	Pro215	Prostate 6	525.59	
	Pro125XB	Prostate 7		556.05
	Pro23B	Prostate 8	1891.4	1118.6
5	ProC280	Prostate 9	454.3	
	Pro20XB	Prostate 10	1332.6	
	Pro34B	Prostate 11		362.91
	Pro65XB	Prostate 12		135.06
	Pro69XB	Prostate 13		179.67
10	Pro10R	Prostate 14 (prostatitis)	143.82	
	Pro20R	Prostate 15 (prostatitis)	397.79	
	Pro258	Prostate 16 (BPH)	216.6	
	Pro263C	Prostate 17 (BPH)	601.25	
	Pro267A	Prostate 18 (BPH)	200.28	
15	Pro271A	Prostate 19 (BPH)	111.43	
	Pro460Z	Prostate 20 (BPH)	53.84	
	ProC032	Prostate 21 (BPH)	56.94	
	SmI21XA	Small Intestine 1	28.8	29.9
	SmIH89	Small Intestine 2	70.8	348.5
20	ClnAC19	Colon 1	22.73	446.47
	ClnAS12	Colon 2	116.97	493.18
	Kid106XD	Kidney 1	86.13	41.14
	Kid107XD	Kidney 2	0.26	35.14
	Lng47XQ	Lung 1	5.13	20.98
25	Lng60XL	Lung 2	13.93	114.78
	Lng75XC	Lung 3	16.47	53.79
	Mam12X	Mammary Gland 1	6.25	10.75
	Mam162X	Mammary Gland 2	1.84	2.54
	Mam42DN	Mammary Gland 3	23.08	35.51

- 30 -

	Ovr10050	Ovary 1	0.9	
	Ovr1028	Ovary 2	261.4	
	Ovr103X	Ovary 3	7	0.1
	Ovr20GA	Ovary 4		0
5	Ovr25GA	Ovary 5		0

0 = Negative

Higher levels of expression were seen in prostate, showing a high degree of tissue specificity for prostate tissue. Of all the analyzed samples different from prostate, only two cancer samples (colon 2 with 116.97 and ovary 2 with 261.4), and 5 normal adjacent tissue samples (small intestine 2, colon 1, colon 2, kidney 1, and lung 2), showed an expression comparable to the mRNA expression in prostate. These results confirmed the tissue specificity results obtained with the panel of normal pooled samples (Table 7).

Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 8 shows higher expression of Prol15 in 3 out of 4 matched prostate cancer tissues (prostate samples 1, 5 & 8).

Altogether, the high level of tissue specificity, plus the higher expression in 75% of the prostate matching samples tested, are indicative of Prol15 being a diagnostic marker for prostate cancer.

Expression of Clone ID 3277219H1 (Prol10):

For the CSG Prol10, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- CGGCAACCTGGTAGTGAGTG - 3' (SEQ ID NO:29)

- 31 -

Reverse Primer

5' - CGCAGCTCCTTGTAACCTTCAG - 3' (SEQ ID NO:30)

The absolute numbers depicted in Table 9 are relative levels of expression of the CSG Prol10 in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 9: Relative Levels of CSG Prol10 Expression in Pooled Samples

10

15

20

Tissue	NORMAL
Brain	6.61
Heart	0.7
Kidney	0.74
Liver	7.94
Lung	11.88
Mammary	22.78
Muscle	6.77
Prostate	3.01
Small Intestine	1
Testis	2.58
Thymus	13.74
Uterus	2.61

The relative levels of expression in Table 9 show that Prol10 mRNA expression is not as high in normal prostate (3.01) compared with all the other normal tissues analyzed.

The absolute numbers in Table 9 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 10.

The absolute numbers depicted in Table 10 are relative levels of expression of Prol10 in 33 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from

- 32 -

the normal adjacent sample for that same tissue from the same individual.

Table 10: Relative Levels of CSG Prol10 Expression in Individual Samples

5	Sample ID	Tissue	Cancer	Matching Normal Adjacent
	Pro12B	Prostate 1	11.8	0.3
	Pro78XB	Prostate 2	14.3	6.3
	Pro101XB	Prostate 3	33.2	10.7
	Pro13XB	Prostate 4	0.3	0.4
10	Pro23XB	Prostate 5	25.5	14.4
	Pro20XB	Prostate 6	43.3	4
	Pro34XB	Prostate 7	31.8	18.7
	Pro65XB	Prostate 8	26.9	3.4
	Pro69XB	Prostate 9	12.5	7
15	Lng75XC	Lung 1	1.9	3
	Lng90X	Lung 2	5.5	0.5
	LngAC11	Lung 3	9.3	9.7
	LngAC32	Lung 4	11.2	2.2
	Lng47XQ	Lung 5	11.3	0.3
20	Lng60XL	Lung 6	29.1	6.8
	Mam12B	Mammary Gland 1	19.8	0
	Mam603X	Mammary Gland 2	13.7	0
	Mam82XI	Mammary Gland 3	73.5	0
	MamA04	Mammary Gland 4	0	24.6
25	MamB011X	Mammary Gland 5	17.4	2
	MamC012	Mammary Gland 6	0	12.8
	MamC034	Mammary Gland 7	0	61
	Mam12X	Mammary Gland 8	14	2.2
	Mam59X	Mammary Gland 9	33	2.2

- 33 -

	MamA06X	Mammary Gland 10	16.4	0.8
	Liv15XA	Liver 1	4.7	0.6
	Liv42X	Liver 2	7.5	2.6
	Liv94XA	Liver 3	0.4	1.4
5	ClnAS43	Colon 1	52.9	1.4
	ClnAS45	Colon 2	2.1	0.8
	ClnAS46	Colon 3	39.8	3.7
	SmI21X	Small Intestine 1	0.9	0.1
	SmIH89	Small Intestine 2	5.8	0.9

10 0 = Negative

The levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 10 shows overexpression of Prol10 in 8 of the 9 primary prostate cancer tissues compared with their respective normal adjacent (except prostate 4). Thus, there was overexpression in 88.88% of the cancer prostate tissue as compared to the prostate matching samples tested (total of 9 prostate matching samples).

Although not tissue specific, Prol10 mRNA expression is upregulated in prostate cancer tissues. The mRNA overexpression in 88.88% of the primary prostate matching cancer samples tested is indicative of Prol10 being a diagnostic marker for prostate cancer. Prol10 also showed overexpression in several other cancers tested including small intestine, colon, liver, mammary and lung (see Table 10). Accordingly Prol10 may be a diagnostic marker for other types of cancer as well.

- 34 -

Expression of Clone ID 1857415; Gene ID 346880 (Prol13):

For the CSG Prol13, real-time quantitative PCR was performed using the following primers:

Forward Primer

5' - CGGGAACCTACCAGCCTATG - 3' (SEQ ID NO:31)

Reverse Primer

5' - CAGGCAACAGGGAGTCATGT - 3' (SEQ ID NO:32)

The absolute numbers depicted in Table 11 are relative levels of expression of the CSG Prol13 in 12 normal different tissues. All the values are compared to normal thymus (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 11: Relative Levels of CSG Prol13 Expression in Pooled Samples

Tissue	NORMAL
Brain	0.03
Heart	0
Kidney	0.01
Liver	0
Lung	0
Mammary Gland	0
Muscle	0.04
Prostate	489.44
Small Intestine	0.02
Testis	0.35
Thymus	1
Uterus	0.13

The relative levels of expression in Table 11 show that Prol13 mRNA expression is higher (489.44) in prostate compared with all the other normal tissues analyzed. Testis, with a relative expression level of 0.35, uterus (0.13), thymus (1.0), kidney (0.01) and brain (0.03) were among the other tissues expressing lower mRNA levels for Prol13. These results establish that Prol13 mRNA expression is highly specific for prostate.

- 35 -

The absolute numbers in Table 11 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 12.

The absolute numbers depicted in Table 12 are relative levels of expression of Prol13 in 78 pairs of matching and 25 unmatched tissue samples. All the values are compared to normal thymus (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual. In cancers (for example, ovary) where it was not possible to obtain normal adjacent samples from the same individual, samples from a different normal individual were analyzed.

Table 12: Relative Levels of CSG Prol13 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matched or Unmatched Normal Adjacent
Pro780B/781B	Prostate 1	375.58	446.29
Pro1291B/1292B	Prostate 2	1060	31
Pro139B96/140B96	Prostate 3	41	32
Pro209B96/210B96	Prostate 4	505	255
Pro1256B/1257B	Prostate 5	165.79	141.63
Pro1293B/1294B	Prostate 6	1613.7	874.61
Pro694B/695B	Prostate 7	458.6	142.21
Pro1012B/1013B	Prostate 8	1520	864
Pro1222B/1223B	Prostate 9	939	530
Pro845B/846B	Prostate 10	1552.4	374.6
Pro1094B/1095B	Prostate 11	278.37	135.89
Pro650B/651B	Prostate 12	532.81	640.85

- 36 -

	Pro902B/903B	Prostate 13	609.05	415.86
	Pro916B/917B	Prostate 14	699.42	401.24
	Pro9821110A/110B	Prostate 15	156	487.8
	ProS9821326A/26B	Prostate 16	744.4	472.8
5	Pro9407c215	Prostate 17	1389.2	
	Pro9407c234	Prostate 18	305.5	
	Pro9407c280A	Prostate 19	894.5	
	Pro9409C010R	Prostate 20 (prostatitis)	269.7	
	Pro9404C120R	Prostate 21 (prostatitis)	299.2	
10	Pro1000258	Prostate 22 (BPH)	149.6	
	Pro4001263C	Prostate 23 (BPH)	576	
	Pro4001267A	Prostate 24 (BPH)	132.1	
	Pro9411C032	Prostate 25 (BPH)	118.2	
	Pro4001460Z	Prostate 26 (BPH)	276.3	
15	Pro4001271A	Prostate 27 (BPH)	58.7	
	Kid1064D/65D	Kidney 1	0	0.1
	Kid1079D/1080D	Kidney 2	0.3	0.02
	Kid1097D/1098D	Kidney 3	35.14	0.32
	Kid1024D/1025D	Kidney 4	1.31	0
20	Kid1183D/1184D	Kidney 5	24.79	0
	Kid1242D/1243D	Kidney 6	0	0
	Bld469K	Bladder 1		2.88
	Bld467K/468K	Bladder 2	2.65	
	Bld327K/328K	Bladder 3	0	4.05
25	Bld470K	Bladder 4		1.64
	Bld665T/664T	Bladder 5	0.21	1.99

- 37 -

	Bld1496K/1497K	Bladder 6	13.55	1.14
	Bld1721K/1722K	Bladder 7	120.16	1.34
	Tst239X/240X	Testis 1	31.5	0.73
	TstS9820647A/47B	Testis 2	15.7	0
5	TstS9820663A/663B	Testis 3	72	1.4
	SknS9821248A/248B	Skin 1	1.8	0.5
	SknS99448A/448B	Skin 2	251.6	0
	Skn99816A/816B	Skin 3	33	0.7
	Sto4004864A4/B4	Stomach 1	14.12	0
10	Sto4004509A3/B1	Stomach 2	40.74	39
	SmI9807A212A/213A	Small Intestine 1	0.1	0
	SmI9802H008/H009	Small Intestine 2	5.8	0.1
	Cln9608B012/B011	Colon 1	4.5	0
	Cln9709c074ra/073ra	Colon 2	65.8	3.1
15	Cln4004709A1/709B1	Colon 3	1.1	0.9
	Cln9405C199/C200	Colon 4	34.76	0.73
	Cln9707c004gb/006ga	Colon 5	90.26	0.96
	Cln96-09-B004/B003	Colon 6	17.9	20.64
	Cln9612B006/B005	Colon 7	17.56	0.3
20	Cln9705F002D/F001C	Colon 8	21.39	0
	ClnCXGA	Colon 9	429.14	142.69
	Pan10343a	Pancreas 1	0	0
	Pan776P/777P	Pancreas 2	0	0.15
	Pan9210/9220	Pancreas 3	7.36	0
25	Pan714L/715L	Pancreas 4	13.57	0.11
	Pan824P/825P	Pancreas 5	0	0
	Lng476Q/477Q	Lung 1	0	0
	Lng605L/606L	Lung 2	0	0.1
	Lng11145B/11145C	Lung 3	85.9	0

- 38 -

	Lng0008632A/32B	Lung 4	23.85	0
	Lng750C/751C	Lung 5	0.32	0.25
	Lng8890A/8890B	Lung 6	10.63	0
	Lng8926A/8926B	Lung 7	15.37	0
5	Lng0010239A/39B	Lung 8	26.17	0
	Lng9502C109R/110R	Lung 9	0.68	0
	LngS9821944a/44b	Lung 10	0	0
	Mam00042D01/42N01	Mammary Gland 1	8.5	0
	Mam59XC	Mammary Gland 2	61.07	0
10	Mam9706A066G/67C	Mammary Gland 3	4.84	0
	Mam14153a1C	Mammary Gland 4	9.72	6.99
	Mam1620F/1621F	Mammary Gland 5	0.91	0
	Mam00014D05	Mammary Gland 6	2.45	0
	End10479B/D	Endometrium 1	133.43	1.12
15	End9705A125A/126A	Endometrium 2	0	0.39
	End9704C281A/282A	Endometrium 3	23.5	1.56
	End680o97/681o97	Endometrium 4	88.89	79.02
	Utr13590/13580	Uterus 1	0.2	0
	Utr850U/851U	Uterus 2	0	0
20	Utr14170/14180	Uterus 3	14	0.4
	Utr233U96/234U96	Uterus 4	8.65	4.64
	CvxVNM00052D01/52N01	Cervix 1	0.82	77.15
	CvxVNM00083D01/83N01	Cervix 2	0.78	221.48
	CvxND00023D01/23N01	Cervix 3	3.25	15.22
25	Ovr10370/10380	Ovary 1	0.1	0
	Ovr10050	Ovary 2	18.96	
	Ovr1028	Ovary 3	0	
	Ovr14638A1C	Ovary 4	3.2	
	Ovr14603A1D	Ovary 5	882.3	
30	Ovr7730	Ovary 6	0	

- 39 -

	Ovr9702C018GA	Ovary 7		0.15
	Ovr206I	Ovary 8		0
	Ovr9702C020GA	Ovary 9		0
	Ovr9702C025GA	Ovary 10		0
5	Ovr9701C035GA	Ovary 11		0.07
	Ovr9701C050GB	Ovary 12		0.58

0 = Negative

In the analysis of matching samples, the higher levels of expression were in prostate, showing a high degree of tissue specificity for prostate tissue. In addition to the higher expression levels in prostate cancer samples, Prol13 expression was found to be either induced (where not expressed in normal adjacent tissues) or somewhat upregulated in several other cancers. However, the relative expression and the fold increase in prostate cancer samples far exceeds that in other cancer tissues and is highly significant.

Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 12 shows overexpression of Prol13 in 13 out of 16 primary prostate cancer tissues compared with their respective normal adjacent (prostate samples 2, 3, 4, 5, 6 7, 8, 9, 10, 11, 13, 14, 16). Thus, there was overexpression in the cancer tissue for 81.25% of the prostate matching samples tested. The median for the level of expression in prostate cancer tissue samples is 609, whereas the median for all other cancers is only 7.93, with the exception of one colon sample, colon 9, whose expression was similar to that found in prostate cancer tissues.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 81.25% of the primary prostate matching samples tested are indicative of Prol13 being a

- 40 -

diagnostic marker for prostate cancer. Expression was also found to be higher in other cancer tissues compared with their respective normal adjacent tissues (kidney, bladder, testis, skin, stomach, small intestine, colon, pancreas, lung, mammary, endometrium, uterus, and ovary) thus indicating Prol13 to be a pan cancer marker.

Expression of Clone ID 1810463H1 (Prol14):

For the CSG Prol14, real-time quantitative PCR was performed using the following primers:

10 Forward Primer

5'- TGGGCATCTGGGTGTCAA - 3' (SEQ ID NO:33)

Reverse Primer

5'- CGGCTGCGATGAGGAAGTA - 3' (SEQ ID NO:34)

The absolute numbers depicted in Table 13 are relative levels of expression of the CSG Prol14 in 12 normal different tissues. All the values are compared to normal muscle (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

20 **Table 13: Relative Levels of CSG Prol14 Expression in Pooled Samples**

Tissue	NORMAL
Brain	9.7
Heart	0.7
Kidney	414.4
Liver	4
Lung	882.2
Mammary	44
Muscle	1
Prostate	1951
Small Intestine	22
Testis	367.1
Thymus	25.8
Uterus	139.6

35 The relative levels of expression in Table 13 show that Prol14 mRNA expression is higher (1951) in prostate compared with all the other normal tissues analyzed. Lung, with a relative

- 41 -

expression level of 882.2, kidney 414.4, testis 367.1 and uterus 139.6, are the other tissues expressing higher levels of mRNA for Prol14. These results establish Prol14 mRNA expression to be more specific for prostate than other tissues examined.

The high level of tissue specificity is indicative of Prol14 being a diagnostic marker for diseases of the prostate, especially cancer.

Expression of Clone ID zr65g11 (Prol18):

For the CSG Prol18, real-time quantitative PCR was performed using the following primers:

Forward Primer

5'- GCCCATCTCCTGCTTCTTTAGT - 3' (SEQ ID NO:35)

Reverse Primer

5'- CGTGGAGATGGCTCTGATGTA - 3' (SEQ ID NO:36)

The absolute numbers depicted in Table 14 are relative levels of expression of the CSG Prol18 in 12 normal different tissues. All the values are compared to normal kidney (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 14: Relative Levels of CSG Prol18 Expression in Pooled Samples

Tissue	NORMAL
Colon	0.87
Endometrium	19282
Kidney	1
Liver	0
Ovary	86.22
Pancreas	0
Prostate	962.1
Small Intestine	0
Spleen	0.75
Stomach	0.54
Testis	343.7
Uterus	1064

- 42 -

The relative levels of expression in Table 14 show that Prol18 mRNA expression is the 3rd highest in prostate (962.1) next to endometrium (19282) and uterus (1064), which are female-specific tissues. Testis, with a relative expression level of 343.7 is the only other male tissue expressing moderate levels of mRNA for Prol18. These results establish Prol18 mRNA expression to be highly specific for reproductive tissues including the prostate.

The absolute numbers in Table 14 were obtained analyzing pools of samples of a particular tissue from different individuals. They cannot be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 15.

The absolute numbers depicted in Table 15 are relative levels of expression of Prol18 in 59 pairs of matching and 21 unmatched samples. All the values are compared to normal kidney (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 15: Relative Levels of CSG Prol18 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Pro12B	Prostate 1	41700.7	22242.83
ProC234	Prostate 2	40087	
Pro78XB	Prostate 3	4075.6	7066.7
Pro109XB	Prostate 4	334.4	777.2
Pro84XB	Prostate 5	11684	58290
Pro101XB	Prostate 6	21474.13	100720.8
Pro91X	Prostate 7	14849	33717
Pro13XB	Prostate 8	202.57	146.91

- 43 -

	ProC215	Prostate 9	73243	
	Pro125XB	Prostate 10	629.6	521.4
	Pro23B	Prostate 11	157532.6	110654.4
	Pro90XB	Prostate 12	2317	64134
5	ProC280	Prostate 13	42020	
	Pro20XB	Prostate 14	2909.31	
	Pro34B	Prostate 15	29610	23264
	Pro110	Prostate 16	13354	30991
	Pro65XB	Prostate 17	10126	11270
10	Pro69XB	Prostate 18		2671.42
	Pro326	Prostate 19	9962.3	19231
	Pro10R	Prostate 20 (prostatitis)	27355	
	Pro20R	Prostate 21 (prostatitis)	21081	
	Pro258	Prostate 22 (BPH)	79916.32	
15	Pro263C	Prostate 23 (BPH)	108924.5	
	Pro267A	Prostate 24 (BPH)	92910.22	
	Pro271A	Prostate 25 (BPH)	57004.4	
	Pro460Z	Prostate 26 (BPH)	57449.23	
	ProC032	Prostate 27 (BPH)	45781.44	
20	Kid106XD	Kidney 1	3.08	217.36
	Kid107XD	Kidney 2	0	38.36
	Kid109XD	Kidney 3	0	123.5
	Kid10XD	Kidney 4	17.69	67.8
	Kid11XD	Kidney 5	16.74	360.8
25	Kid124D	Kidney 6	0	167.4
	Bld32XK	Bladder 1	0	0
	Bld47K	Bladder 2		36.38
	Bld66X	Bladder 3	0	4.52
	BldTR14	Bladder 4	0	12.17

- 44 -

	BldTR17	Bladder 5	0	0
	Bld46XK	Bladder 6	16.5	0
	Tst39X	Testis 1	116.6	24.35
	Tst647T	Testis 2	856.16	43.5
5	StoAC44	Stomach 1	0	0
	StoAC93	Stomach 2	0	0
	SmI21XA	Small Intestine 1	68.45	0
	SmIH89	Small Intestine 2	0	0
	ClnAC19	Colon 1	149	21.33
10	ClnAS12	Colon 2	0	0
	ClnB34	Colon 3	0	0
	ClnB56	Colon 4	13.04	5.22
	ClnAS43	Colon 5	0	0
	Lng47XQ	Lung 1	0	0
15	Lng60XL	Lung 2	0	0
	Lng75XC	Lung 3	0	3.38
	Lng90X	Lung 4	0	0
	LngBR26	Lung 5	0	26.82
	Pan10343	Pancreas 1	50.47	0
20	Pan77X	Pancreas 2	281.1	0
	Pan92X	Pancreas 3	18.41	0
	Pan71XL	Pancreas 4	0	0
	Pan82XP	Pancreas 5	0	0
	PanC044	Pancreas 6	0	0
25	Mam12X	Mammary Gland 1	0	0
	Mam162X	Mammary Gland 2	0	0
	Mam42DN	Mammary Gland 3	0	0
	MamS127	Mammary Gland 4	12.58	0
	Mam14DN	Mammary Gland 5	0	0
30	End28XA	Endometrium 1	331.9	1824

- 45 -

	End3AX	Endometrium 2	27825	65839
	End4XA	Endometrium 3	10.3	15935
	Utr1410	Uterus 1	18885	18116
	Utr23XU	Uterus 2	3358	7674
5	CvxKS52	Cervix 1	0	0
	CvxKS83	Cervix 2	0	0
	Ovr10050	Ovary 1	72.86	
	Ovr1028	Ovary 2	0	
	Ovr638A	Ovary 3	0	
10	Ovr63A	Ovary 4	90.88	
	Ovr7730	Ovary 5	1.21	
	Ovr10400	Ovary 6	5.08	
	Ovr1050	Ovary 7	0	
	Ovr1118	Ovary 8	7.41	
15	Ovr103X	Ovary 9		32.78
	Ovr20GA	Ovary 10		0
	Ovr25GA	Ovary 11		1173.83
	Ovr35GA	Ovary 12		313.4
	Ovr50GB	Ovary 13		823.1
20	Ovr18GA	Ovary 14		40.6
	Ovr206I	Ovary 15		1264
	Ovr230A	Ovary 16		1285

0 = Negative

In the analysis of matching samples, the higher levels of expression were in prostate, endometrium, testis, and ovary showing a high degree of tissue specificity for reproductive tissues. These results confirmed the tissue specificity results obtained with the panel of normal pooled samples (Table 14).

Furthermore, the levels of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an

- 46 -

indication of specificity for the cancer (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 15 shows overexpression of Prol18 in 5 out of 14 primary prostate cancer tissues (prostate samples 1, 8, 5 10, 11, 15) compared with their respective normal adjacent. Thus, there was overexpression in the cancer tissue for 35.71% of the prostate matching samples tested (total of 14 prostate matching samples). Expression of Prol18 was similarly higher in 3 unmatched cancer tissues (prostate samples 9, 13, 14), 10 2 prostatitis (prostate samples 20, 21), and 6 benign hyperplasia tissues (prostate samples 22 through 27).

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 35.71% of the primary prostate matching samples tested are indicative of Prol18 being a 15 diagnostic marker for prostate cancer.

- 47 -

What is claimed is:

1. A method for diagnosing the presence of prostate cancer in a patient comprising:

(a) determining levels of CSG in cells, tissues or bodily fluids in a patient; and

(b) comparing the determined levels of CSG with levels of CSG in cells, tissues or bodily fluids from a normal human control, wherein a change in determined levels of CSG in said patient versus normal human control is associated with the presence of prostate cancer.

2. A method of diagnosing metastases of prostate cancer in a patient comprising:

(a) identifying a patient having prostate cancer that is not known to have metastasized;

(b) determining CSG levels in a sample of cells, tissues, or bodily fluid from said patient; and

(c) comparing the determined CSG levels with levels of CSG in cells, tissue, or bodily fluid of a normal human control, wherein an increase in determined CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

3. A method of staging prostate cancer in a patient having prostate cancer comprising:

(a) identifying a patient having prostate cancer;

(b) determining CSG levels in a sample of cells, tissue, or bodily fluid from said patient; and

(c) comparing determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in determined CSG levels in said patient versus the normal human control is associated with a cancer which is progressing and a decrease in the determined CSG levels is associated with a cancer which is regressing or in remission.

- 48 -

4. A method of monitoring prostate cancer in a patient for the onset of metastasis comprising:

- (a) identifying a patient having prostate cancer that is not known to have metastasized;
- 5 (b) periodically determining levels of CSG in samples of cells, tissues, or bodily fluid from said patient; and
- (c) comparing the periodically determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal human control, wherein an increase in any one of the
- 10 periodically determined CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

5. A method of monitoring a change in stage of prostate cancer in a patient comprising:

- 15 (a) identifying a patient having prostate cancer;
- (b) periodically determining levels of CSG in cells, tissues, or bodily fluid from said patient; and
- (c) comparing the periodically determined CSG levels with levels of CSG in cells, tissues, or bodily fluid of a normal
- 20 human control, wherein an increase in any one of the periodically determined CSG levels in the patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease is associated with a cancer which is regressing in stage or in remission.

25 6. A method of identifying potential therapeutic agents for use in imaging and treating prostate cancer comprising screening molecules for an ability to bind to CSG wherein the ability of a molecule to bind to CSG is indicative of the molecule being useful in imaging and treating prostate cancer.

30 7. The method of claim 1, 2, 3, 4, 5 or 6 wherein the CSG comprises SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12,

- 49 -

13, 14, 15, 16, 17, 18, 19 or 20 or a polypeptide encoded thereby.

8. An antibody which specifically binds CSG.

9. A method of imaging prostate cancer in a patient
5 comprising administering to the patient an antibody of claim 8.

10. The method of claim 9 wherein said antibody is labeled with paramagnetic ions or a radioisotope.

11. A method of treating prostate cancer in a patient
10 comprising administering to the patient an antibody of claim 7.

12. The method of claim 11 wherein the antibody is conjugated to a cytotoxic agent.

SEQUENCE LISTING

<110> Salceda, Susana
 Recipon, Herve
 Cafferkey, Robert
 diaDexus, LLC

<120> Method of Diagnosing, Monitoring, Staging, Imaging and
 Treating Prostate Cancer

<130> DEX-0052

<140>

<141>

<150> 60/104,737

<151> 1998-10-19

<160> 36

<170> PatentIn Ver. 2.0

<210> 1

<211> 188

<212> DNA

<213> Homo sapiens

<400> 1

```

ggtaaacacc tgcttttatc atcagaacaa agaggctgtg tcccctgccc tatgagggtcc 60
atcttctgaga gttgtggcta atgggcaaga aggttggggc tttagagatt tgggataaaag 120
atatcaaaca ccagaaagggt agaaagaagt gatcagatta gggttactta ggtgatgata 180
tgaactct
188

```

<210> 2

<211> 9819

<212> DNA

<213> Homo sapiens

<400> 2

```

cagctggggg ctaccaggt ccatgtcttg gacatgttga gagtttttct ggaaggcagg 60
gatacagtgt ggtccaaaaa cacacaaatg cccctactgg cccaggggtt gtcacaatag 120
actggaaggg tgacacatcc caggcgcttg ccaccatca cacgcacctc ctaccactg 180
gcatccttcc accccaggca cacacaaagc ctcagtccag agatcaactc tggactcagc 240
tctgaatttg catatcctgt gtgtagattc attcttcata acctctgccc agcctagctt 300
gtgtatcatt tttttttctc tattagggga ggagcccgtc ctggcactcc cattggcctg 360
tagattcacc tcccctgggc agggccccag gaccaggat aatatctgtg cctcctgccc 420
agaaccctcc aagcagacac aatggtaaga atggtgcctg tcctgctgtc tctgctgctg 480
cttctgggtc ctgctgtccc ccaggagaac caagatggtg agtggggaaa gcaagggatg 540

```

ggtgctggag aggactggaa ggaggtgagg aacaggacat gtggctggga gacaggctgg 600
 atgcagctgg gataccctgg cataccggcag gaatgggtgc ccaaggctgt caactccctc 660
 agctcacaca cttccaggag cattcaggga gcctctgcgc tggcccga aa taagaccttc 720
 aggaatctga atctaaaacc cctagtttac agtgaaaaca aagactccaa agaccaagcg 780
 acctgcttgg ggtagacagt caggacggag taggaacct atgcctggag ctgcttctgc 840
 tcctgttcct tccctccttc cgatggctgg gtacacctgc ctgacgctga ggaaaagaga 900
 gagcagcccc aaggggaaag tgggaaggca ggttggctgg agggatggtg ctagaaggaa 960
 acccgtgccc aaatcccaca ctcagacacc actgcagtgg gtctggaagg cgagtggctg 1020
 gaagagaaga gagtgggagc tccgggagat caagagtcac tcctaggata agggaaggag 1080
 gctgtttgtg gcatgagaat gtgcaggata aagacatgga agcgaatggc ttctcagttg 1140
 tgtgagttta aaattcatga catttacaaa ttgtcagaaa aggtgttata tgtttgttat 1200
 ataacaatca ctttggaatg ttaatctgat tctgtgccaa aatctgaatt actcagggtt 1260
 ctccagagaa acagaactaa taggtggtac acatatacat atatatgtac gtacacatac 1320
 atacatacac tgtatacaca tggatacaca cacacatagg aagagattta catatatgta 1380
 tacaaaagag agagagagta gagattttatt ttaagaaatt gactcacact attgggagga 1440
 gtaacaagtc ctaaactctc agagccggcc agcaggctgg agaccaggg aagagttgat 1500
 gtcttagtct tgattccaag ggcagactgt aggcagaatt ctttcctctt taggggacat 1560
 ctgaggcttt ttctcttaag gccttcaact gattggatga agcccaccac tatggagagt 1620
 aatccacttt actcaaggtc tactgatttt tttgtaaatt aaaaaaaaaa ctgtgggtgc 1680
 atagtatgtg tatatattta tgggggtacat gagaggtttt gattcaggca tgcaatgtga 1740
 aataatcaca tcatcaaaaa tgaggtatcc atcccttcaa gcttttatcg tttgtgttac 1800
 agacaatcca attatacttt tttggttatt ttagttttta aaagtatttg attattttatt 1860
 tattttatta tttttgagac agagtctcac tctgtcacc aggcaggagt gcagtggcat 1920
 gatctcggct cactgcaacc tccgcctccc aggttcaagc aattttcctg cctcagctctc 1980
 ctgagtagct aggactacag gcacctgcca ccacacctgg ctaatttttt tgtattttta 2040
 gtagagacgg tttcatcatg ttggccaggc tagtcttgat atcctgacct cgtgatctgc 2100
 ccgccttggc tccccaaagt gccgggatta caggtgtcag caactgcgc tggcctctct 2160
 tttggttatt taaaagtgtg caattaaatt atgattatta ttattatttt tgagatggat 2220
 tcttgttctg tcaccaggc tggagtgcag tggcgtgatc ttggcttact gcaaacctcc 2280
 gcctgttggg ttcaagcaat tatcttgct cgggtgtaca ctgccacaca cggctaactt 2340
 atgtattttt aatagagata gggcttcacc atgttggcta gactggctct gacctcttga 2400
 cctcaagtga tccactcact tcagcctccc agagtgtgg aattacaggc acgagccacc 2460
 acacctggcc ccagttaaatt tattattgac tatagtcacc ctgttgtgct atcaaatagt 2520
 aggtcttatt cattcttctt tttttttttt tttttgtgac agagttgccc aggttggaat 2580
 gcagtgggtg aatcttggct cactgcaacc tctgcctccc gggcttaagc gattctcctg 2640
 cctcagcctt ctgagtcgct gggactacag gtgtgtgcca ccacgcccgg ctaattttatg 2700
 tatttttagt agagatgggg tttcaccatg ttggccaggc tggtttcgaa ctctgacct 2760
 caagtgacct acctgcctca gcttcccaaa gtgttggaat tacaggcatg agccaccaca 2820
 cctggcccca gttaaattat tattcactgg agtcactttg ttgtgctatc aaatagtttt 2880
 ctaactattt tttttgtacc cattaaccac cctcccaatt tcccccaac cctgccacta 2940
 cccttccag cctttggtaa ccatccttct actctctatg tccatgaatt caattgtagg 3000
 gtctactgat ttaaaggcta atcacattta gacactcagg agcaagaata attttagtaa 3060
 ttgaactagg attctgcat atgacctcca acatcattag cacctgtgta aattgtatca 3120
 taaaataatt atggaactat tatggaaatg tccctctctc ccagatccca ccttgtacca 3180
 aaatgcaagg tacaacccc ggaattctga gctccatcct agtcttacc tgtgctaatt 3240
 cagtctgggt catttcttga atttcttgg aaattctcct ttctaccctt tctaactata 3300
 tgtatttgc aggttaagc agaagtgtta atttttttt tttttgagat ggagccttgc 3360
 tttgtcacct aggtgaagt gcagtggcat gatctcagct cactgcaagc tccgcctccc 3420

gggttcatgc cattctcctg cctcagcctc ctgagtagct gggactacag gcacccgcca 3480
 ccatgcttgg ctaatttttt gaattcttag tagagacggg gtttcacat gttagccagg 3540
 atgggtctcga tctcctgacc tcgtgatcca ccgcctcgg cccctaaag tgctgggatt 3600
 acaggcgtga gccactgagc ccggacgaaa tgtaatttg ttttttttga gacggagtct 3660
 cactctgtca tccaagctgg agtgacgtgg catgatcttg gcttggttga acctctgcct 3720
 ctctgggtca agtgattttc ctgcctcagc ctccagcatg actgggatta caggccccga 3780
 ccaccatgcc cagctaattt ttgtattttt taatagagat ggggtttcac catgttgccc 3840
 aggctggctc tcaactcctg atctcaagta atctgcctgc cttggcctcc caaagtccctg 3900
 ggattacagg catgagccac ggagcccagc ctagaaatgt taattttctaa cgcagtgcag 3960
 attccatgca cactgggcaa ggttccattc ctccatgggg tgactcaggg atccaggcca 4020
 attgcatatt gagactcttt catattatcc tgtggccttc aaagtcgtca cctctaggga 4080
 tgagaacaaa aagggaagc cagctggtag ggtcttgga aagaagaaag acatcacttc 4140
 tgctcacatt ctcttttgac aaaactcagt cacatggtcc caatatactc tcgagggtggc 4200
 tgagtaatgt tatcttctta tgtgtcaagc agaggaaata atgtagtga gacacaggat 4260
 ggtctctgaa atatcatctc aggcattgaaa gtagagcata ttcacttgag tgagcctcca 4320
 gtggtgtgaa gttgatggca ggagaaagag ctggggaaga aaaggccagt ggcaggcttc 4380
 cctcctagc cctatgcagc ccacagtggt gacccttgca tggacctca ccatcagaat 4440
 ctcttctttt gcaggctgtt actctctgac ctatatctac actgggctgt ccaagcatgt 4500
 tgaagacgtc ccgcgtttc aggccttggt ctactcaat gacctcagt tcttttagata 4560
 caacagtaaa gacaggaagt ctacagccat gggactcttg agacaggtgg aaggaatgga 4620
 ggattggaag caggacagcc aacttcagaa ggccaggag gacatcttta tggagacct 4680
 gaaagacatt gtggagtatt acaacgacag taacggtcag tgaataacag accacagggg 4740
 tggaggtct aaccaagag gcagccccc cagtgtgagt ggcaaggat cagcaggatg 4800
 gaaatagtcc caatcccagg ggaagaacag gagacacagc agaaacacag acatgtccgc 4860
 atcccacca cccacagca caggtgctcc ccgttcccc atcaattgcc ccatcctcat 4920
 ccaggcctc aggtcacaca ggaagtgtg gcagagtcac ttcctatcca ggcacctatg 4980
 acctctcacc tccacacccc acccatcgga ggctgatacc ccgtgagaa ggcacagac 5040
 tcaccctgt ccaggaggt tgcttgaga gtgagccact ctcaaagtc ctcagacctg 5100
 ggctcacctg gtggtctgc cagtcctagc tgttgacagt gaaacgttcc caaatatct 5160
 ggttgaaatc tgcaaacatt ggagcactga gacctacct caaacaagtc tgtaatat 5220
 aactatgtct gttctatgaa ggatgtcaca gtctgtcctg atctcccttg cagctccatc 5280
 acctagcaca ggttacagcc aatattggct caattgaaat ttgtggaatc cacagagaaa 5340
 agcaccggc acacaccgta gcccatgctg ggggtcagg aagtgtgga ttcaaaactg 5400
 tgggtgtta gatttcttg gagccctaaa gttctcctt accatacgt gcagaccag 5460
 gaagggccac ctgcgtatg gtcagaggag ctggtggcag agcccggtgca gagatggctc 5520
 ctgtgcccc ggccagtgct tctttctcct aaaccacact gccagcccca aggcagccaa 5580
 cctcaggtct ggtgaactgc tgggtttaa ttatcataga gtgggtgtca aaagatgggc 5640
 tactaagtac aaaaatgcc aagtgctac atgggatctg aagattttca aaaggaggca 5700
 agaaagagat aggcagatgt ttcaaggatg tgggtgggg gaggtcttg taaggaaaat 5760
 ggcccaggct gtgtgtcagc aataggagag gagggggcac aggtgatcag aaaagacact 5820
 gggggaagca ttgatggaca ggaatagaaa tggcaaagt gataattaag aggaaggagg 5880
 atgaggagat gaacacaggg tattagaaaa taatagaagg cagggttggtg tggctcactc 5940
 ttgtaatccc agcacttttg gaggtgagg caggcagatc acctagggtc aggagtctga 6000
 gaccagccc gccaacatgg tgaaccctg tctctactaa taatacaaaa atagcctggc 6060
 atggtggcac acgtctgtg tccagctac tcaggaggct gaggcaggag aattgcttga 6120
 acccaggagg cagaggttac agtggccaaa atcctacat tgcactacag cctgggtgac 6180
 aagagtgaac cgttgtctaa aaacaaaaa caaaaaacaa aaaaaggaaa taatagtagc 6240
 tgacatttac tgagcactta ctttgtgcca ggccatcta tgagcatata taatgctcag 6300

aatagcccc taaaacagtg ctcttggeat tgccatttca gaggtgagga aatagaggca 6360
 caggagattg agtggctcca gttcaggcaa cacaccaggt gggggtgggg ggctggggag 6420
 agacctggga cgtgagccca gacagcttga gagctttcag agtctatgcc aacagcacca 6480
 accagtgtg ggtaaacacc tgcttttate atcagaacaa agaggctgtg tccctgccc 6540
 tatgaggtcc atttctgaga gttgtggcta atgggcaaga aggttggggc tttgagatt 6600
 tgggataaag atatcaaaca ccagaaaggt agaaagaagt gatcagatta gggttactta 6660
 ggtgatgata tgaactcttc ctagaactga gagaaaaaga gagccttcct ttactcatat 6720
 gaaatcacaa ataatttcta tccaatttgg aagtacactt tgggtgagtt gtgacagctt 6780
 cctcaggact cagcataaat tcaaacaaat aattgtcctt agaagagatg ctatagaaga 6840
 gatagaaata tattcatatt ctgtagcttt tttttttttg agatggagtt ttgctctgt 6900
 caccgaagct ggagtgcagt gatgcaatct cagctcactg caaactttgc ctctgggtt 6960
 caagggattc tcctgcctca gcctcccgat aactgggact acaggctaca ggcatgtgtc 7020
 actactcctg gtttaattttt tttttttttt ttttaagactg agtcttgctc tgtctttcag 7080
 gctgatgtac aatggctcca tctcggtcca ctacaacttc tgtccccag gttcaagcga 7140
 ttctcctgcc tcagcctcat gagtagctgg gattacaggc atgtgccagc acaccagca 7200
 aatttttgta ttttttagtag agatgaggtc ttaccatgtt ggccaggctg gtctcaaact 7260
 cctgacctca ggtgatcctt tggectcagc ctccctaact gctgggatta caggcatgag 7320
 ccactgcgtc cagcctaatt ttatatTTTT ggtagagatg gggtttcacc atattggcca 7380
 ggctggcttc gaactcatga cctaagggtga tccatcctcc tcagccttc aaagtgtctg 7440
 gattacaagt gtgagccact gggcctgggt cttttttttt tttttttttt tttttttttt 7500
 tgagataggg tctcactctg tcaccaggc tgaaatgcag tagtgtgatt ttggctcatt 7560
 gcagccttga ctccccaggc tgaagtgatc ctccacctc agcctcctga gtagctgggg 7620
 ctacaggcat gcaccacat gctgcgctaa tttttatatt tttttagtg gtgggatttc 7680
 gccatatcac cctggctgggt ctggaacccc tgggctcaag cgatccactc gcttcagctt 7740
 ctcaaagtgc tgggattaca ggcattgagc acagcgccca ggctgtagct ctcttaagga 7800
 ggaacatatc tcatctgaga caaacctgaa atgccaaacc aaactgagtt agccctctc 7860
 tgtctgttgt atatattgga gtaataacct atttgtcttg ataaagggat tgcatgcttg 7920
 aattgcaaaa acctttattt cttttgggtt gcccaatgtg caagactaag agttattttg 7980
 ataaatttct caccaggctg actgtctctc tgtggggctg ggggagttt cagggtctca 8040
 cgtattgcag ggaaggtttg gttgtgagat cgagaataac agaagcagcg gagcattctg 8100
 gaaatattac tatgatggaa aggactacat tgaattcaac aaagaaatcc cagcctgggt 8160
 ccccttcgac ccagcagccc agataacca gcagaagtgg gaggcagaac cagtctacgt 8220
 gcagcggggc aaggcttacc tggaggagga gtgccctgcg actctgcgga aatacctgaa 8280
 atacagcaaa aatatcctgg accggcaagg tactcactgc ttcctgctcc ccagtactga 8340
 gccagaata aaagacgatc tcaggctagg agctcaggca acatcttagt ccggtctcat 8400
 ctgttcctgg atgtccctca gaccccccag tttcatcttt taggatttat tccttcctg 8460
 ggataatata atttgtggtc caaaaagaac atcatcaaaa tttcaggcag aatgggcccag 8520
 gaaggccatt ctttcttgat gagtgtcccc aaatcatctc caattaacag acaaggagct 8580
 tgaggttagg gaggtgaggg taacactgtc tgtaagaggc agagctggga ctcaaattcc 8640
 agatttcaga ttccaaatcc catcgttttt tatctctaca atgatgcctc ccatctgggt 8700
 ggtggagaga agggaggcgt gtaaaagtca gcccagaag gacaagagca agccagtgtg 8760
 agcggaattg atggctgcaa gctgagactt ggattggaga cgtagtgaga ctcaggattg 8820
 tgcagtgtg caggggaagt gttgctggat agaggcatgg gctgaaccaa gcagctggac 8880
 tgagactggg ggacagaact ccaaagccca ctgagatgtg ggaaaacatg gagaagcaca 8940
 cggagcatte acaacttatt gccgtcagag tcaatacatg ggtgaggtgg ggattgggca 9000
 agagggaaa cgtcagcctt ccctgatatt ctggaaagtc tcccggggct ggggggtggc 9060
 aggtacagag ctctcagctc tgctgatcgc tgacatccag ggggtgggggt aggaagagac 9120
 ctgggcccgg agaagtccac ctcaagcctg cagtgtcaca ctctatccct ccacagatcc 9180

```

tccctctgtg gtggtcacca gccaccaggc cccaggagaa aagaagaaac tgaagtgcct 9240
ggcctacgac ttctaccag ggaaaattga tgtgcactgg actcggggccg gcgaggtgca 9300
ggagcctgag ttacggggag atgttcttca caatggaaat ggcaacttacc agtcctgggt 9360
ggtggtggca gtgccccgc aggacacagc cccctactcc tgccacgtgc agcacagcag 9420
cctggcccag cccctcgtgg tgccctggga ggccagctag gaagcaaggg ttggaggcaa 9480
tgtgggatct cagacccagt agctgccctt cctgcctgat gtgggagctg aaccacagaa 9540
atcacagtca atggatccac aaggcctgag gagcagtgtg gggggacaga caggaggtgg 9600
atttgagac cgaagactgg gatgcctgtc ttgagtagac ttggacccaa aaaatcatct 9660
caccttgagc ccacccccac cccattgtct aatctgtaga agctaataaa taatcatccc 9720
tccttgcccta gcataacaga gaatcctttt tttaacgggtg atgcgctgta gaaatgtgac 9780
tagattttct cattggttct gccctcaagc actgaattc 9819

```

<210> 3

<211> 250

<212> DNA

<213> Homo sapiens

<400> 3

```

cgccccctgcg ccgccgagcc agctgccaga atgccgaact ggggaggagg caagaaatgt 60
gggggtgtgtc agaagacggt ttactttgcc gaagagggtc agtgccaagg caacagcttc 120
cataaatcct gcttcctgtg catggtctgc aagaagaatc tggacagtac cactgtggcc 180
gtgcatggtg aggagattta ctgcaagtcc tgctacggca agaagtatgg gcccaaaggc 240
tatggctacg 250

```

<210> 4

<211> 1900

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (16)

<220>

<221> unsure

<222> (18)

<220>

<221> unsure

<222> (20)

<220>

<221> unsure

<222> (1887)

<220>

<221> unsure

<222> (1894)

<400> 4

```

acgccttccg cggagnanan caaaacggcg cgcaggccgg gcgcacccag ccgccacttc 60
cgagagcgcc tgccgcccct ggcgcccgcg agccagctgc cagaatgccg aactggggag 120
gaggcaagaa atgtgggggtg tgtcaagaag acggtttact ttgccgaaga ggttcagtgc 180
gaaggcaaca gcttccataa atcctgcttc ctgtgcatgg tctgcaagaa gaatctggac 240
agtaccactg tgggccgtgc atggtgagga gatttactgg caagtccctg ctacggcaag 300
aagtatgggc ccaaaggcta tggctacggg ccagggcgca ggcaccctca gcactgacaa 360
gggggagtcg ctgggtatca agcacgagga agcccctggg ccacaggccc accaccaacc 420
ccaatggcat ccaaatttgc ccagaagatt ggtggtcccg agcgctgccc ccgatgcagc 480
caggcagctc atgctgcgga gaaggtgatt ggtgctggga agtcctggca taaggcctgc 540
tttcgatgtg ccaagtgtgg caaaggcctt gagtcaacca ccctgggcag acaaggatgg 600
cgagatttac tgcaaaggat gttatgctaa aaacttcggg cccaagggct ttggttttgg 660
gcaaggagct ggggccttgg tccactctga gtgaggccac catcaccac cacaccctgc 720
ccactcctgc gcttttcac gccattccat tcccagcagc tttggagacc tccaggatta 780
tttctctgtc agccctgcc catatcacta atgacttgaa cttgggcac tggctccctt 840
tgggttgggg gtctgcctga ggtccacccc cactaaaggg ctccccaggc ctgggatctg 900
acaccatcac cagtaggaga cctcagtggt ttgggtctag gtgagagcag gcccctctcc 960
ccacacctcg cccacagag ctctgttctt agcctcctgt gctgctgtc catcatcagc 1020
tgaccaagac acctgaggac acatcttggc acccagagga gcagcagcaa caggctggag 1080
ggagagggaa gcaagaccaa gatgaggagg ggggaaggct gggttttttg gatctcagag 1140
attctcctct gtgggaaaga ggttgagctt cctggtgtcc ctgagagtaa gcctgaggag 1200
tcccagctta gggagtccac tattggaggc agagaggcat gcaggcaggg tcctaggagc 1260
ccctgcttct ccaggcctct tgcccttgag tctttgtgga atggatagcc tcccactagg 1320
actgggagga gaataaccca ggtcttaagg accccaaagt caggatgttg tttgatcttc 1380
tcaaacatct agttccctgc ttgatgggag gatcctaag aaatacctga aacatatatt 1440
ggcatttatc aatggctcaa atcttcattt atctctggcc ttaaccctgg ctctgaggc 1500
tgcgccagc agagcccagg ccagggtctt gttcttgcca cacctgcttg atcctcagat 1560
gtggaggagg gtaggcactg cctcagctt catccaaaca cttttccct tgccctgaga 1620
cctcagaatc ttccttttaa cccaagaccc tgccctcttc actccacct tctccaggga 1680
cccttagatc acatcactcc acccctgcc gggcccagg taggaatagt ggtgggagga 1740
aggggaaagg gctgggcctc accgctccca gcaactgaaa ggacaacact atctggagcc 1800
accactgaa agggctgcag gcatgggctg tacccaagct gatttctcat ctggtcaata 1860
aagctgttta gaccagaaaa aaaaaanaaa aaanaaaagg 1900

```

<210> 5

<211> 273

<212> DNA

<213> Homo sapiens

<400> 5

```

gatgcatcaa aagagctgca agttctccac attgacttct tgaatcagga caacgccgtt 60
tctcaccaca catgggagtt ccaaacgagc agtcctgtgt tccggcgagg acagggtgtt 120
cacctgcggc tgggtctgaa ccagccccta caatcctacc accaactgaa actggaattc 180
agcacagggc cgaatcctag catcgccaaa cacaccctgg tgggtgctga cccgaggacg 240
ccctcagacc actacaactg gcaggcaacc ctt 273

```

<210> 6

<211> 3021

<212> DNA

<213> Homo sapiens

<400> 6

tgtggaagca ccaggcatca gagatagagt cttccctggc attgcaggag agaattctgaa 60
gggatgatgg atgcatcaaa agagctgcaa gttctccaca ttgacttctt gaatcaggac 120
aacgccgttt ctcaccacac atgggagttc caaacgagca gtcctgtgtt ccggcgagga 180
cagggtgtttc acctgcggct ggtgctgaac cagcccctac aatcctacca ccaactgaaa 240
ctggaattca gcacagggcc gaatcctagc atcgccaaac acaccctggt ggtgctcgac 300
ccgaggacgc cctcagacca ctacaactgg caggcaaccc ttcaaaatga gtctggcaaa 360
gaggtcacag tggctgtcac cagttcccc aatgccatcc tgggcaagta ccaactaaac 420
gtgaaaactg gaaaccacat ccttaagtct gaagaaaaca tcctatacct tctcttcaac 480
ccatggtgta aagaggacat ggttttcatg cctgatgagg acgagcgcaa agagtacatc 540
ctcaatgaca cgggctgcca ttacgtgggg gctgccagaa gtatcaaatg caaaccttgg 600
aactttgggtc agtttgagaa aaatgtcctg gactgctgca tttccctgct gactgagagc 660
tccctcaagc ccacagatag gagggacccc gtgctggtgt gcaggggccat gtgtgctatg 720
atgagctttg agaaaggcca gggcgtgctc attgggaatt ggactgggga ctatgaaggt 780
ggcacagccc catacaagtg gacaggcagt gccccgatcc tgcagcagta ctacaacacg 840
aagcaggctg tgtgctttgg ccagtgtctg gtgtttgctg ggatcctgac tacagtgtctg 900
agagcgttgg gcatcccagc acgcagtgtg acaggcttcg attcagctca cgacacagaa 960
aggaacctca cgggtggacac ctatgtgaat gagaatggca agaaaatcac cagtatgacc 1020
cacgactctg tctggaattt ccatgtgtgg acggatgcct ggatgaagcg accggatctg 1080
cccaagggtc acgacggctg gcaggctgtg gacgcaacgc cgcaggagcg aagccagggt 1140
gtcttctgct gtgggcatc accactgacc gccatccgca aagggtgacat ctttattgtc 1200
tatgacacca gattcgtctt ctcaagaagt aatgggtgaca ggctcatctg gttggtgaag 1260
atgggtgaatg ggcaggagga gttacacgta atttcaatgg agaccacaag catcgggaaa 1320
aacatcagca ccaaggcagt gggccaagac aggcggagag atatcaccta tgagtacaag 1380
tatccagaag gtcctctga ggagaggcag gttcatggat catgccttc tccttctcag 1440
ttctgagagg gagcacagac gacctgtaaa agagaacttt cttcacatgt cggtaacaatc 1500
agatgatgtg ctgctgggaa actctgttaa tttcacctg attcttaaaa ggaagaccgc 1560
tgccctacag aatgtcaaca tcttgggctc ctttgaacta cagttgtaca ctggcaagaa 1620
gatggcaaaa ctgtgtgacc tcaataagac ctgcagatc caaggtaag tatcagaagt 1680
gactctgacc ttggactcca agacctacat caacagcctg gctatattag atgatgagcc 1740
agttatcaga ggtttcatca ttgcggaaat tgtggagtct aaggaaatca tggcctctga 1800
agtattcacg tctttccagt accctgagtt ctctatagag ttgcctaaca caggcagaat 1860
tggccagcta cttgtctgca attgtatctt caagaatacc ctggccatcc ccttgactga 1920
cgtcaagttc tctttggaaa gcctgggcat ctctcacta cagacctctg accatgggtg 1980
agtctgcctg aggacggtgc agcctggtga gaccatccaa tcccaaataa aatgcacccc 2040
aataaaaatg gacccaagaa atttatcgtc aagttaagtt ccaaacaagt gaaagagatt 2100
aatgtctaga agattgttct catcaccaag tagccttgct tgatgctgtg gagccttagt 2160
tgagatttca gcatttccta ccttgtggct tagctttcag attatggatg attaaatttg 2220
atgacttata tgagggcaga ttcaagagcc agcagggtcaa aaaggccaac acaaccataa 2280
gcagccagac ccacaaggcc aggtcctgtg ctatcacagg gtcaccttct tttacagtta 2340
gaaacaccag ccgaggccac agaatcccat ccctttcctg agtcatggcc tcaaaaatca 2400
gggcccacat tgtctcaatt caaatccata gatttcgaag ccacagattc tctccctgga 2460
gcaagcatga ctatgggcag ccagtgctg ccacctgctg acgacccttg agaagctgcc 2520
atatcttcag gccatgggtt caccagccct gaaggcacct gtcaactgga gtgtctcttc 2580

```

agcactggga tgggcctgat agaagtgc at tctcctccta ttgcctccat tctcctctct 2640
ctatccctga aatccaggaa gtccctctcc tgggtgtcca agcagtttga agcccaatct 2700
gcaaggacat ttctcaaggg ccatgtggtt ttgcagacaa cctgtcctc aggcctgaac 2760
tcaccataga gacccatgtc agcaaacggg gaccagcaaa tctctctccc ttattctaaa 2820
gctgcccctt ggggagactcc agggagaagg cattgtctcc tccctgggtg gaactctttc 2880
tttgggtatt catccactat cctggcaact caaggctgct tctgttaact gaagcctgct 2940
ccttcttggt ctgccctcca gagatttgct caaatgatca ataagcttta aattaaactc 3000
tacttcaaga aaaaaaac g 3021

```

<210> 7

<211> 267

<212> DNA

<213> Homo sapiens

<400> 7

```

gaacattcca gatacctatc attactcgat gctgttgata acagcaagat ggctttgaac 60
tcagggtcac caccagctat tggaccttac tatgaaaacc atggatacca accggaaaac 120
ccctatcccc cacagcccac tgtgggtccc actgtctacg aggtgcatcc ggctcagtac 180
taccggtccc ccggtgcccc gtacgccccg agggctcctga cgcaggcttc caaccccgctc 240
gtctgcacgc agcccaaata cccatcc 267

```

<210> 8

<211> 3443

<212> DNA

<213> Homo sapiens

<400> 8

```

gggaggggcg ggccgagtag gcgcgagcta agcaggaggc ggaggcggag gcggagggcg 60
agggggcgggg agcgccgcct ggagcgcggc aggtcatatt gaacattcca gatacctatc 120
attactcgat gctgttgata acagcaagat ggctttgaac tcagggtcac caccagctat 180
tggaccttac tatgaaaacc atggatacca accggaaaac ccctatcccc cacagcccac 240
tgtgggtccc actgtctacg aggtgcatcc ggctcagtac taccggtccc ccggtgcccc 300
gtacgccccg agggctcctga cgcaggcttc caaccccgctc gtctgcacgc agcccaaata 360
cccatccggg acagtgtgca cctcaaagac taagaaagca ctgtgcatca ccttgacctc 420
ggggaccttc ctctgtggag ctgcgtggc cgctggccta ctctggaagt tcatgggcag 480
caagtgtccc aactctggga tagagtgcga ctctcagggt acctgcatca acccctctaa 540
ctgggtgtgat ggcgtgtcac actgccccg cggggaggac gagaatcggg gtgttcgcct 600
ctacggacca aacttcattc tttaggtgta ctcatctcag aggaagtcct ggcacctgt 660
gtgccaagac gactggaacg agaactacgg gcggggcgcc tgcagggaca tgggctataa 720
gaataatatt tactctagcc aaggaatagt ggatgacagc ggatccacca gctttatgaa 780
actgaacaca agtgccggca atgtcgatat ctataaaaa ctgtaccaca gtgatgcctg 840
ttcttcaaaa gcagtgggtt ctttacgctg tatagcctgc ggggtcaact tgaactcaag 900
ccgccagagc aggatcgtag gcggcgagag cgcgctccc ggggcctggc cctgggcagg 960
tcagcctgca cgtccagaac gtccacgtgt gcggaggctc catcatcacc ccgagtgga 1020
tcgtgacagc cgccactgac gtggaaaaac ctcttaacaa tccatggcat tggacggcat 1080
ttgcggggat tttgagacaa tctttcatgt tctatggagc cgataccaa gtagaaaaag 1140
tgatttctca tccaaattat gactccaaga ccaagaacaa tgacattgag ctgatgaagc 1200
tgcagaagcc tctgactttc aacgacctag tgaaaccagt gtgtctgccc aaccagggca 1260

```

tgatgctgca gccagaacag ctctgctgga tttccgggtg gggggccacc gaggagaaaag 1320
 ggaagacctc agaagtgctg aacgctgcca aggtgcttct cattgagaca cagagatgca 1380
 acagcagata tgtctatgac aacctgatca caccagccat gatctgtgcc ggcttcctgc 1440
 aggggaacgt cgattcctgc cagggtgaca gtggagggcc tctggctact tcgaagaaca 1500
 atatctgggtg gctgataggg gatacaagct ggggttctgg ctgtgccaaa gcttacagac 1560
 caggagtgtg cggaatgtg atggatttca cggactggat ttatcgacaa atgagggcag 1620
 acggctaata cacatggtct tcgtccttga cgtcgtttta caagaaaaca atgggctgg 1680
 ttttgcttcc ccgtgcatga tttactctta gagatgattc agaggctact tcatttttat 1740
 taacacagtga acttgctggt ctttgccact ctctgccatt ctgtgcaggc tgcagtggct 1800
 cccctgcccc gcctgctctc cctaaccctt tgtccgcaag gggatgatggc cggctgggtg 1860
 tgggcactgg cggtaagtg tggaggagag gggatggaggc tgccccattg agatcttctt 1920
 gctgagtcct ttccaggggc caattttgga tgagcatgga gctgtcacct ctcagctgct 1980
 ggatgacttg agatgaaaaa ggagagacat ggaaagggag acagccaggc ggcacctgca 2040
 gcggctgcct ctggggccac ttggtagtgt cccagccta cctctccaca aggggatttt 2100
 gctgatgggt tcttagagcc ttagcagccc tggatgggtg ccagaaataa agggaccagc 2160
 ccttcatggg tggtagcgtg gtatgcacct tgtaagggga acagaaacat ttttgttctt 2220
 atgggggtgag aatatagaca gtgcccttgg gtgcgagggg agcaattgaa aaggaacttg 2280
 cccctgagcac tcctgggtgca ggtctccacc tgcacattgg gtggggctcc tgggagggag 2340
 actcagcctt cctctctatc ctccttgacc ctgctcctag caccctggag agtgcacatg 2400
 ccccttggtc ctgggcaggg gcgccaagtc tggcaccatg ttggcctctt caggcctgct 2460
 agtcaactgga aattgaggtc catgggggaa atcaaggatg ctgagtttaa ggtacactgt 2520
 ttccatgtta tgtttctaca cattgctacc tcagtgtctc tggaaactta gcttttgatg 2580
 tctccaagta gtccaccttc atttaactct ttgaaactgt atcatctttg ccaagtaaga 2640
 gtggtggcct atttcagctg ctttgacaaa atgactggct cctgacttaa cgttctataa 2700
 atgaatgtgc tgaagcaaag tgcccatggt ggcggcgaag aagagaaaaga tgtgttttgt 2760
 tttggactct ctgtggtccc ttccaatgct gtgggtttcc aaccagggga agggctccctt 2820
 ttgcattgcc aagtgcata accatgagca ctactctacc atggttctgc ctcctggcca 2880
 agcaggctgg tttgcaagaa tgaaatgaat gattctacag ctaggactta accttgaaat 2940
 ggaaagtctt gcaatcccat ttgcaggatc cgtctgtgca catgcctctg tagagagcag 3000
 cattcccagg gaccttgga acagttggca ctgtaagggt cttgctcccc aagacacatc 3060
 ctaaaagggt ttgtaatggt gaaaacgtct tccttcttta ttgccccttc ttatttatgt 3120
 gaacaactgt ttgtcttttt ttgtatcttt ttttaaactgt aaagttcaat tgtgaaaatg 3180
 aatatcatgc aataaatta tgcgattttt ttttcaaagt aaccactgca tctttgaagt 3240
 tctgcctggt gagtaggacc agcctccatt tccttataag ggggtgatgt tgaggtgct 3300
 ggtcagagga ccaaagggtg ggcaaggcca gacttggtgc tcctgtggtt ggtgccctca 3360
 gttcctgcag cctgtcctgt tggagaggtc cctcaaatga ctccttctta ttattctatt 3420
 agtctgtttc catgggcgtg ata 3443

<210> 9

<211> 254

<212> DNA

<213> Homo sapiens

<400> 9

gtgctgcacc aggccacat cctgccaag actgggacag tgccctgga ggtacggctc 60
 ctggaggcct cccgtgcctt cgaggtgtca gagaacggca acctggtagt gagtgggaag 120
 gtgtaccagt gggatgaccc tgacccagg ctcttcgacc acccgaaaag cccaccccc 180
 aacccacgg agccctctt cctggcccag gctgaagttt acaaggagct gcgtctgct 240

ggctacgact acgg

254

<210> 10

<211> 8470

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (4131)

<220>

<221> unsure

<222> (5117)

<220>

<221> unsure

<222> (5552)

<400> 10

cgcccgctcga cacggcagcg gccccggcct cctctctcgc cgcgcttcag cctcccgcctc 60
 cgcccgctc cagcctcgct ctccgcccgc cgcaccgccc cccgcccct caccagagca 120
 gccatggagg aggtggtgat tgccggcatg tccgggaagc tgccagagtc ggagaacttg 180
 caggagtctt gggacaacct catcgccggt gtggacatgg tcacggacga tgaccgtcgc 240
 tggaaggcgg ggctctacgg cctgccccgg cggctccggca agctgaagga cctgtctagg 300
 tttgatgcct ccttcttcgg agtccacccc aagcaggcac acacgatgga cctcagctg 360
 cggctgctgc tggaagtcac ctatgaagcc atcgtggacg gaggcataa cccagattca 420
 ctccgaggaa cacacactgg cgtctgggtg ggcgtgagcg gctctgagac ctccgaggcc 480
 ctgagccgag accccgagac actcgtgggc tacagcatgg tgggctgcca gcgagcgatg 540
 atggccaacc ggctctcctt cttcttcgac ttcagagggc ccagcatcgc actggacaca 600
 gcctgctcct ccagcctgat ggccctgcag aacgcctacc aggccatcca cagcgggcag 660
 tgccctgccg ccacgtggg gggcatcaat gtccctgctga agcccaacac ctccgtgcag 720
 ttcttgaggc tggggatgct cagccccgag ggcacctgca aggccttcga cacagcgggg 780
 aatgggtact gccgctcgga ggggtgtggt gccgtcctgc tgaccaagaa gtccctggcc 840
 cggcgggtgt acgccaccat cctgaacgcc ggcaccaata cagatggctt caaggagcaa 900
 ggcgtgacct tccccacag ggatatccag gagcagctca tccgctcgtt gtaccagtcg 960
 gccggagtgg cccctgagtc atttgaatac atcgaagccc acggcacagg caccaagggtg 1020
 ggcgaccccc aggagctgaa tggcatcacc cgagccctgt gcgccacccg ccaggagccg 1080
 ctgctcatcg gctccaccaa gtccaacatg gggcaccgag agccagcctc ggggctggca 1140
 gccctggcca aggtgctgct gtccctggag caggggctct gggcccccac cctgcacttc 1200
 catagcccca accctgagat ccagcgtg ttggatgggc ggctgcaggt ggtggaccag 1260
 cccctgcccc tccgtggcgg caacgtgggc atcaactcct ttggcttcgg gggctccaaa 1320
 cgtgcacatc atcctgaggc ccaacacgca gccgcccccc gcacccggcc cacatgccac 1380
 cctgccccgt ctgctgcggg ccagcggacg caccctgag gccgtgcaga agctgctgga 1440
 gcagggcctc cggcacagcc agggcctggc ttccctgagc atgtgaacga catcgccggt 1500
 gtccccgacc accgccatgc cttccgtgg ctacgctgtg ctgggtggtg agacgcgggtg 1560
 gccagaggt gcagcaggtg cccgctggcg agcgcgccgt ctggttcac tgctctggga 1620
 tgggcacaca gtggcgcggg atggggctga gcctcatgcg cctggaccgc ttccgagatt 1680

ccatcctacg ctccgatgag gctgtgaacc gattcggcct gaaggtgtca cagctgctgc 1740
 tgagcacaga cgagagcacc tttgatgaca tcgtccattc gtttgtgagc ctgactgcca 1800
 tccagatagg cctcatagac ctgctgagct gcatggggct gaggccagat ggcacgtcg 1860
 gccactccct gggggaggtg gcctgtggct acgccgacgg ctgcctgtcc caggaggagg 1920
 ccgtcctcgc tgcctactgg aggggacagt gcatcaaaga agcccatctc ccgcccggcg 1980
 ccatggcagc cgtgggcttg tcctgggagg agtgtaaaca gcgctgcccc ccggcggtgg 2040
 tgccccgcgc cacaactcca aggacacagt caccatctcg ggacctcagg ccccggtgtt 2100
 tgagttcgtg gagcagctga ggaaggaggg tgtgtttgcc aaggaggtgc ggaccggcgg 2160
 tatggccttc cactcctact tcatggaggc catcgacccc ccactgctgc aggagctcaa 2220
 gaaggtgatc cgggagccga agccacgttc agcccgctgg ctacgacct ctatccccga 2280
 ggcccagtg gacagcagcc tggcacgcac gtccctccgc gactacaatg tcaacaacct 2340
 ggtgagccct gtgtgttcc aggaggccct gtggcacgtg cctgagcacg cgggtgtgt 2400
 ggagatcgcg cccacgccc tgcctgaggc tgcctgaag cgtggcctga agccgagctg 2460
 caccatcctc cccctgatga agaaggatca cagggacaac ctggagtctc tcctggccgg 2520
 catcggcagg ctgcacctct caggcatcga cgccaacccc aatgccttgt tcccacctgt 2580
 ggagtcccca gctccccgag gaactccct catctcccca ctcatcaagt gggaccacag 2640
 cctggcctgg gacgcgccgg ccgcccaggga ctccccaac ggttcagggt cccctcagc 2700
 caccatctac acatgcacac caagctccga gtctcctgac cgctacctgg tggaccacac 2760
 catcgacggt cgcgtcctct tccccgccac tggctacctg agcatagtgt ggaagacgct 2820
 ggcccagacc ctgggcctgg gcgtcgagca gctgcctgtg gtgtttgagg atgtggtgt 2880
 gcaccaggcc accatcctgc ccaagactgg gacagtgtcc ctggagggtac ggctcctgga 2940
 ggctcctcgt gccttcgagg tgtcagagaa cggcaacctg gtagtgagt ggaaggtgta 3000
 ccagtgggat gacctgacc ccaggctctt cgaccacccc gaaagcccca ccccaaccc 3060
 caggagccc ctcttctgag cccaggctga agtttacaag gagctgcgtc tgcgtggcta 3120
 cgactacggc cctcatcttc agggcatcct ggaggccagc ctggaagggt actcggggag 3180
 gctgctgtgg aaggataatg ggtgagttca tggacaccat gctgcagatg tccatcctgg 3240
 gtcggccaag caggcctgt acctgcccac ccgtgtcacc gccatccaca tcgacctgc 3300
 caccacagg cagaagctgt acacactgca ggacaaggcc caagtggctg acgtggtgt 3360
 gagcaggtgg ctgagggtca cagtggccgg aggcgtccac atctccgggc tccacactga 3420
 gtcggccccg cggcggcagc aggagcagca ggtgcccatc ctggagaagt tttgcttcac 3480
 tccccacag gaggaggggt gcctgtctga gcacgtgccc ctcgaggagg agctgcaact 3540
 gtgcaagggg ctggtcgagg cactcgagac caaggtgacc cagcaggggc tgaagatggt 3600
 ggtgcccgga ctggatgggg ccagatccc cccgggaccc ctcacagcag gaactgcccc 3660
 ggctgttgct ggctgcctgc aggttcacg tcaacgggaa cctgcagctg gagctggcgc 3720
 aggtgctggc ccaggagagg cccaagctgc cagaggaccc tctgctcagc ggctcctgg 3780
 actccccggc actcaaggcc tgcttgga caagcgtgga gaacatgccc agcctgaaga 3840
 tgaaggtggt ggaggtgctg gccggccacg gtcacctgta ttcccgcatc ccaggcctgc 3900
 tcagccccca tccctgctg cagctgagct acacggccac cgaccgccac cccaggccc 3960
 tggaggctgc ccaggccgag ctgcagcagc acgacgtg ccagggccag tgggatccc 4020
 cagacctgc cccagcgcc ctgggcagcg cggacctcct ggtgtgcaac tgtgctgtg 4080
 ctgcccctcg ggacccgct cagctctcag caacatggtg gctgccctga nagaagggg 4140
 ctttctgctc ctgcacacac tgctccgggg gcacccctc ggggacatcg tggccttct 4200
 cactccact gagccgcagt atggccaggg catcctgagc caggacgct gggagagcct 4260
 cttctccagg gtgtcgctgc gcctggtgg cctgaagaag tccttctacg gctccacgct 4320
 cttcctgtgc cgccggccca ccccgaggga cagcccatc ttctgcccg tggacgatac 4380
 cagcttccgc tgggtggagt ctctgaagg catcctggct gacgaagact ctttcccgcc 4440
 ctgtgtggct gaaggccatc aactgtcca cctcgggcgt ggtgggcttg gtgaactgtc 4500
 tccgcccaga gcccgcgga acgctccggt gtgtgctgct ctccaacctc agcagacct 4560

cccacgtccc ggaggtggac ccgggctccg cagaactgca gaaggtgttg cagggagacc 4620
 tgggtgatgaa cgtctaccgc gacggggcct ggggggcttt ccgccacttc ctgctggagg 4680
 aggacaagcc tgaggagccg acggcacatg cctttgtgag caccctcacc cggggggacc 4740
 tgtccctcca tccgctgggt ctgctcctcg ctgcccctg cccagcccac ctgcccctggc 4800
 gcccagctct gcacggtcta ctacgcctcc ctcaacttcc gcgacatcat gctggccact 4860
 ggcaagctgt cccctgatgc catcccaggg aagtggacct cccaggacag cctgctaggt 4920
 atggagtctt cgggcccaga cgccagcggc aagcgtgtga tgggactggg gcctgccaa 4980
 ggccctggcca cctctgtcct gctgtcaccg gacttctctt gggatgtgcc ttccaactgg 5040
 acgctggagg agggggcctc ggtgcctgtc gtctacagca cggcctacta cgcgctgggt 5100
 gtgctggggc ggggtgcncct cggggagacg ctgctcatcc actcgggctc gggcgggctg 5160
 ggccaggccg ccacgcccac cgccctcagt ctgggctgcc gcgtcttcac caccgtgggg 5220
 tcggctgaga agcgggcgta cctccaggcc aggttcccc agctcgacag caccagcttc 5280
 gccaaactccc gggacacatc cttcgagcag catgtgctgt ggcacacggg cgggaagggg 5340
 gttgacctgg tcttgaactc cttggcgga gagaaagtgc aggccagcgt gaggtgcttg 5400
 gctacgcacg gtcgcttctt ggaaattggc aaattcgacc tttctcagaa ccaccgctc 5460
 ggcattggcta tcttcttgaa gaacgtgaca ttccacgggg tctactgga tgcgttcttc 5520
 aacgagagca gtgctgactg gcgggagggt tnggcgcttg tgcaggccgg catccgggat 5580
 ggggtggtag gggccctcaa gtgcacgggt ttccatgggg cccagggtga ggacgccttc 5640
 cgctacatgg ccaaggga gcacattggc aaagtctcg tgcagggtgct tgcggaggag 5700
 ccggaggcag tggctgaagg gggccaaacc caagctgatg tcggccatct ccaagacctt 5760
 ctgcccggcc cacaagagct acatcatcgc tgggtggtctg ggtggcttcg gcctggagtt 5820
 ggcgagtggt ctgatacagc gtgggggtga gaagctcgtg ttgacttctc gctccgggat 5880
 ccggacaggc taccaggcca agcaggctcc cggtggagg cgccaggcgg tacagggtga 5940
 ggtgtccacc agcaacatca gctcactgga gggggcccg ggcctcattg ccgaggcggc 6000
 gcagcttgag gcccgtaggc ggcgtcttca acctggcgt ggtcttgaga gatggcttgc 6060
 tggagaacca gacccagag ttcttccagg acgtctgcaa gccaagtac agcggcacc 6120
 tgaacctgga cagggtagc cgagggcgtg ccctgagctg gactactttg tggctcttctc 6180
 ctctgtgagc tgcgggctg gcaatgcggg acagagcaac tacggctttg ccaatttccg 6240
 ccatggagcg tatctgtgag aaacgcccgc acgaaggcct cccaggcctg gccgtgcagt 6300
 ggggcccatt cggcgacgtg ggcatttttg tggagacgat gagaccaac gacacgatcg 6360
 tcagtggcac gctgccccag cgcattggct cctgcctgga ggtgctggac ctcttctga 6420
 accagcccca catggtcctg agcagctttg tgctggctga gaaggctgcg gcctataggg 6480
 acagggacag ccagcgggac ctgggtggagg ccgtggcaca catcctgggc atccgcgact 6540
 tggctgctgt caacctggac agctcactgg cggacctggg cctggactcg ctcatgagcg 6600
 tggaggtgag ccagacgctg gagcgtgagc tcaacctggg gctgtccgtg cgcgaggtgc 6660
 ggcaactcac gctccggaat ctgcaggagc tgcctcaaaa ggcggtatgag gccagcgagc 6720
 tgggcatgcc ccacgcccga ggaggatggg ctggcccagc agcagactca gctgaacctg 6780
 cgctccctgc tgggtgaacct ggaggggccc acctgatgc ggctcaactg ccgtgcagag 6840
 ctcgagcggg cccctgttcc tgggtgaccc aattcgaggg ctccaccacc gtgttccaca 6900
 gcctggcctc ccggctcagc atccccacct atggcctgca gtgcaccga gctgcgcccc 6960
 ttgacagcat ccacagcctg gctgcctact acatcgactg catcaggcag gtgcagcccc 7020
 agggccctta ccgctgggac ggctactcct acggggcctg cgtggccttt gaaatgtgct 7080
 cccagctgca ggcccagcag agcccagccc ccaccacaa cagcctcttc ctgttcgacg 7140
 gctcggccac ctacgtactg gcctacaccc agagctaccg ggcaaagctg accccaggct 7200
 gtgaggctga ggctgagacg gagccatat gcttcttctg gcagcagttc acggacatgg 7260
 agcacaacag ggtgctggag gcgctgctgc cgtgaaggg cctagaggag cgtgtggcag 7320
 ccgcccgtga cctgatcatc aagagccacc agggcctgga ccgccaggag ctgagctttg 7380
 cggcccggtc cttctactac aagctgcgtg ccgctgagca gtacacaccc aaggccaagt 7440

accatggcaa cgtgatgcta ctgcgcgcca agacgggtgg cgcctacggc gaggacctgg 7500
 gcgcggacta caacctctcc caggtatgcg acgggaaagt atccgtccac gtcatcgagg 7560
 gtgaccaccg cacgctgctg gagggcagcg gcctggagtc catcatcagc atcatccaca 7620
 gctccctggc tgagccacgc gtgagcgtgc gggagggcta ggcccggtgc cccgcctgcc 7680
 accggaggtc actccaccat cccaccccca tcccacccca ccccgccat gcaacgggat 7740
 tgaagggtcc tgccgggtggg accctgtccg gccagtgcc actgcccccc gaggctagct 7800
 agacgtaggt gttaggcatg tcccaccac cgcgcgcctc ccacggcacc tcggggacac 7860
 cagagctgcc gacttggaga ctctgtgtct gtgaagagcc ggtggtgccc gtgcccgcag 7920
 gaactggggc tgggcctcgt gcgcccgtgg ggtctgcgct tggctcttct gtgcttggat 7980
 ttgcatatct attgcattgc tggtagagac cccagggcct gtccaccctg ccaagactcc 8040
 tcaggcagcg tgtgggtccc gcaactctgc cccatttccc cgatgtcccc tgcgggcgcg 8100
 ggcagccacc caagcctgct ggctgcggcc ccctctcgcc caggcattgg ctgagcccg 8160
 tgagtggggg gtcgtgggcc agtccccgag gactggggcc ctgcacaggc acacagggcc 8220
 cggccacacc cagcggcccc ccgcacagcc acccgtgggg tgcctgcctt atgcccggcg 8280
 ccgggcacca actccatgtt tgggtgttgt ctgtgttgt ttttcaagaa atgattcaaa 8340
 ttgctgcttg gattttgaaa ttactgtaa ctgtcagtgt acacgtctgg accccgttcc 8400
 atttttacac caatttggtg aaaatgtctg tctcagcctc ccacaattaa accgcatgtg 8460
 atctccaaaa 8470

<210> 11

<211> 812

<212> DNA

<213> Homo sapiens

<400> 11

gccgcagcca atcagcgcgc gtgcccgggc ccctgcgtct cttgcgtcaa gacggccgtg 60
 ctgagcgaat gcaggcgact tgcgagctgg gagegattta aaacgctttg gattcccccg 120
 gcctgggtgg ggagagcgag ctgggtgccc cctagattcc ccgccccgc acctcatgag 180
 ccgacctcgc gctccatgga gcccggaat tatgccacct tggatggagc caaggatata 240
 gaaggcttgc tgggagcggg aggggggcgg aatctggtcg cccactcccc tctgaccagc 300
 cccccagcgc cgcctacgct gatgcctgct gtcaactatg ccccttggga tctgccaggc 360
 tcggcggagc gccaaagcaa tgccaccat gccctggggg gcccagggg acgtcccag 420
 ctcccgtgcc ttatggttac tttggaggcg ggtactactc ctgccgagtg tcccggagct 480
 cgctgaaacc ctgtgcccag gcagccaccc tggccgcgta ccccgcgag actcccacgg 540
 ccggggaaga gtaccccagc cgcgccactg agtttgctt ctatccggga tatccgggaa 600
 cctaccagcc tatggccagt tacctggacg tgtctgtggt gcagactctg ggtgctcctg 660
 gagaaccgcg acatgactcc ctgttgccgt tggacagtta ccagtcttg gctctcgctg 720
 gtggctggaa cagccagatg tgttgccagg gagaacagaa cccaccagg cccttttgga 780
 aggcagcatt tgcagactcc agcgggcagc ac 812

<210> 12

<211> 2385

<212> DNA

<213> Homo sapiens

<400> 12

ataagctggg gtaaagtatt ttcgcagttt ctgccttttag gattttatta gcttctctcc 60
 cccaggccgc agccaatcag cgcgcgtgcc cgggcccctg cgtctcttgc gtcaagacgg 120


```

ccgtgctgag cgaatgcagg cgacttgcca gctgggagcg atttaaaacg ctttggattc 180
ccccggcctg ggtggggaga gcgagctggg tgccccctag attccccgcc cccgcacctc 240
atgagccgac cctcggtctc atggagcccc gcaattatgc caccttggat ggaagccaagg 300
atatcgaagg cttgctggga gctgggagggg ggcggaatct ggtcgccac tccccctga 360
ccagccaccc agcggcgctc acgctgatgc ctgctgtcaa ctatgcccc ttggatctgc 420
caggctcggc ggagccgcca aagcaatgcc acccatgccc tggggtgccc caggggacgt 480
ccccagctcc cgtgccttat ggttactttg gaggcgggta ctactcctgc cgagtgtccc 540
ggagctcgct gaaacctgt gcccaggcag ccacctggc cgcgtacccc gcggagactc 600
ccacggccgg ggaagagtac cccagccgcc ccactgagtt tgccttctat ccgggatata 660
cgggaacctc ccagcctatg gccagttacc tggacgtgtc tgtggtgcag actctgggtg 720
ctcctggaga accgcgacat gactccctgt tgcctgtgga cagttaccag tcttgggtc 780
tcgctggtgg ctggaacagc cagatgtgtt gccagggaga acagaaccca ccaggtccct 840
tttgaaggc agcatttgca gactccagcg ggcagcacc tcctgacgcc tgcgccttcc 900
gtcgcggccg caagaaacgc attccgtaca gcaaggggca gttgcgggag ctggagcggg 960
agtatgcggc taacaagttc atcaccaagg acaagaggcg caagatctcg gcagccacca 1020
gcctctcgga gcgccagatt accatctggt ttcagaaccg ccgggtcaaa gagaagaagg 1080
ttctcgccaa ggtgaagaac agcgtaccc cttaaagat ctccctgcct ggggtggagg 1140
agcgaagtgt ggggtgtcct ggggagacca ggaacctgcc aagcccaggc tggggccaag 1200
gactctgctg agaggccct agagacaaca ccctccag gccactggct gctggactgt 1260
tcctcaggag cggcctgggt acccagtatg tgcagggaga cggaaccca tgtgacagcc 1320
cactccacca gggttcccaa agaacctggc ccagtcataa tcattcatcc tgacagtggc 1380
aataatcacg ataaccagta ctagtgcga tgatcgtag cctcatattt tctatctaga 1440
gctctgtaga gcactttaga aaccgcttcc atgaattgag ctaattatga ataaatttgg 1500
aaggcgatcc ctttgcaggg aagctttctc tcagacccc ttccattaca cctctcacc 1560
tggtaacagc aggaagactg aggagagggg aacgggcaga ttcgttgtgt ggctgtgatg 1620
tccgtttagc attttctca gctgacagct gggtaggtgg acaattgtag aggtgtctc 1680
ttcctccctc cttgtccacc ccatagggtg taccactgg tcttgaagc acccctctt 1740
aatacgtaga ttttctgtc gtgtgaaat gaagccagca ggctgcccct agtcagtcct 1800
tccttcaga gaaaaagaga tttgagaaag tgctgggta attcaccatt aatttcctcc 1860
ccaaactct ctgagtcttc ccttaatatt tctggtggtt ctgaccaaag caggtcatgg 1920
tttgttgagc atttgggac ccagtgaagt agatgtttgt agccttgcat acttagccct 1980
tcccaggcac aaacggagtg gcagagtggg gccaacctg ttttccagc ccacgtagac 2040
agattcacgt gcggaattct ggaagctgga gacagacggg ctctttgcag agccgggact 2100
ctgagagggg catgagggcc tctgcctctg tgttcattct ctgatgtcct gtacctgggc 2160
tcagtccccg gtgggactca tctcctggcc gcgcagcaa gccagcgggt tcgtgctggg 2220
ccttcctgca ccttaggctg ggggtgggg gcctgccggc gcattctcca cgattgagcg 2280
cacaggcctg aagtctggac aaccgcgaga accgaagctc cgagcagcgg gtcggtggcg 2340
agtagtgggg tcggtggcga gcagttggtg gtgggcccgc gccgc 2385

```

<210> 13

<211> 221

<212> DNA

<213> Homo sapiens

<400> 13

```

dsdnrstact tttctgtgtg gtgcagccct gttggcagtg ggcattctggg tgtcaatcga 60
tggggcatcc tttctgaaga tcttcgggcc actgtcgtcc agtgccatgc agtttgtcaa 120
cgtgggctac ttcctcatcg cagccggcgt tgtggtcttt gctcttggtt tcctgggctg 180

```

ctatggtgct aagactgaga gcaagtgtgc cctcgtgacg t

221

<210> 14

<211> 1533

<212> DNA

<213> Homo sapiens

<400> 14

gggcacgcag acattctggg aagccacttg cccaccccct gggctgcttc ttcttgagat 60
 caggaggggc gttgcccagg gctggtgttg ccagggtggag gcctgctgag gcagtggttg 120
 tggggatcgg tctccaggca gcagggggca gcaggggtcaa ggagaggcta actggccacg 180
 ggtggggcca gcaggcgggc agaaggaggc tttaaagcgc ctaccctgcc tgcagggtgag 240
 cagtgggtgtg tgagagccag gccgtccctc tgcctgcccc ctcagtggca acacccggga 300
 gctgttttgt cctttgtgga gcctcagcag ttccctgctt tcagaactca ctgccaaagag 360
 cctgaacag gagccaccat ggcaagtgtt cagcttcatt aagaccatga tgatcctctt 420
 caatttctc atctttctgt gtggtgcagc cctgttgga gtgggcatct ggggtgtcaat 480
 cgatggggca tcctttctga agatcttcgg gccactgtcg tccagtggca tgcagtttgt 540
 caacgtgggc tacttcctca tcgcagccgg cgttgtggtc tttgctcttg gtttcctggg 600
 ctgctatggt gctaagactg agagcaagtg tgcctcgtg acgttcttct tcatcctcct 660
 cctcatcttc attgctgagg ttgcagctgc tgtggtcgcc ttggtgtaca ccacaatggc 720
 tgagcacttc ctgacgttgc tggtagtgcc tgccatcaag aaagattatg gttcccagga 780
 agacttcact caagtgtgga acaccaccat gaaagggctc aagtgtgtgt gcttcaccaa 840
 ctatacggat tttgaggact caccctactt caaagagaac agtgcccttc cccattcttg 900
 ttgcaatgac aacgtcacca acacagccaa tgaaacctgc accaagcaaa aggctcacga 960
 ccaaaaagta gaggttgct tcaatcagct tttgtatgac atccgaacta atgcagtcac 1020
 cgtgggtggg gtggcagctg gaattggggg cctcgagctg gctgccatga ttgtgtccat 1080
 gtatctgtac tgcaatctac aataagtcca cttctgcctc tgccactact gctgccacat 1140
 gggaaactgtg aagaggcacc ctggcaagca gcagtgttg ggggagggga caggatctaa 1200
 caatgtcact tgggccagaa tggacctgcc ctttctgctc cagacttggg gctagatagg 1260
 gaccactcct ttagggcat gcctgacttt ccttccattg gtgggtggat ggggtggggg 1320
 cattccagag cctctaaggt agccagttct gttgccatt cccccagtct attaaacct 1380
 tgatatgcc cctaggccta gtggtgatcc cagtgtctta ctgggggatg agagaaaggc 1440
 attttatagc ctgggcataa gtgaaatcag cagagcctct ggggtgatgt gtagaaggca 1500
 cttcaaatg cataaacctg ttacaatgtt gcc 1533

<210> 15

<211> 472

<212> DNA

<213> Homo sapiens

<400> 15

tcagagaaaa ctcaaacttt attgagagaa ttttcaaatt ttcagtcaca ttttcaatgt 60
 gacatcagcc atgtgtgtag cttcagcttg tcttcttttt aacttatggc tgcccatctc 120
 ctgcttcttt agtcttagca tgcttaggat taggtggagt cttctctttt acatcagagc 180
 catctccacg ctactccga gtcttttcca gatccatttc ctggcaatca cttctacttt 240
 tacgttcttc gatcgagggt gttccttctc tctctgttcc aggttcaata tctgtattgt 300
 cagttggtgg ttctcttgc tgagattcac cgggagccac gaatgcaacc acatcggggg 360
 cctcctgacc atctcctctt cctctggatc ttgatctcac tcgtgcactc atcgtgcaa 420

ctagaagatc gtgaactgaa gaacttgagt cagcagagag cctggcgaag aa 472

<210> 16

<211> 478

<212> DNA

<213> Homo sapiens

<400> 16

cttcattctt cgccaggctc tctgctgact caagttcttc agttcacgat cttctagttg 60
cagcgatgag tgcacgagtg agatcaagat ccagaggaag aggagatggg caggaggctc 120
ccgatgtggg tgcattcgtg gctcccgggtg aatctcagca agaggaacca ccaactgaca 180
atcaggatat tgaacctgga caagagagag aaggaacacc tccgatcgaa gaacgtaaag 240
tagaaggatga ttgccaggaa atggatctgg aaaagactcg gagtgagcgt ggagatggct 300
ctgatgtaaa agagaagact ccacctaatc ctaagcatgc taagactaaa gaagcaggag 360
atgggcagcc ataagttaaa aagaagacaa gctgaagcta cacacatggc tgatgtcaca 420
ttgaaaatgt gactgaaaat ttgaaaattc tctcaataaa gtttgagttt tctctgaa 478

<210> 17

<211> 198

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (191)

<400> 17

cccgtgtac caccacagca tggtctgcgc cggcggaggg caagaccaga aggactcctg 60
caacgggtgac tctggggggc ccctgatctg caacgggtac ttgcagggcc ttgtgtcttt 120
cggaaaagcc ccgtgtggcc aagttggcgt gccagggtgc tacaccaacc tctgcaaatt 180
cactgagtgg nattaagg 198

<210> 18

<211> 465

<212> DNA

<213> Homo sapiens

<400> 18

tggagatgga gtatgtatctt attttcaaaa aataaatcac catcttcgga ccatttgtag 60
actggaacat ttcgagcaat gagtgcgcca caggagcagag tgccctgggtg actccctgat 120
gttcgcgtca cccccagggc caccttggcg ccgcgatgag cctcgcttcc cactcccggc 180
ctccaactcc cttccctcgc agccgccatt caccttctgc tgtttatttg tctgcagagc 240
gcctggacac cggaagaggc gattccctga gcgcctggag ttggagacaa ttccctgggtc 300
agaatttaaa catctttcta aggttaagcgc tgctccaaaa ctcttcgccg cgtggggact 360
ttgcaccagg ggcggttggg aaggaagttg gccctccacg gggttcctggg caaccgcggc 420
ctgttgaaaa aaggttcttg gtcaaataat ttaacttcgg agggag 465

<210> 19

<211> 204
 <212> DNA
 <213> Homo sapiens

<400> 19
 ggcggaaca ggcgcgctg gacctgtacc cctacgacgc cgggacggac agcggttca 60
 ccttctcctc cccaacttc gccaccatcc cgcaggacac ggtgaccgag ataacgtcct 120
 cctctcccag ccacccggcc aactccttct actaccgcg gctgaaggcc ctgcctccca 180
 tcgccagggt gacactggtg cggc 204

<210> 20
 <211> 294
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (287)

<400> 20
 gagatttctc ttcaatggct tcctgtgagc tagagtttga aaatatctta aaatcttgag 60
 ctagagatgg aagtagcttg gacgattttc attatcatgt aaatcgggtc actcaagggg 120
 ccaaccacag ctgggagcca ctgctcaggg gaaggttcat atgggacttt ctactgcca 180
 aggttctata caggatataa aggtgcctca cagtatagat ctggtagcaa agtaagaaga 240
 aacaacact gatctctttc tgccaccct ctgacccttt ggaactnctc tgac 294

<210> 21
 <211> 22
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic

<400> 21
 atcagaacaa agaggctgtg tc 22

<210> 22
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic

<400> 22
 atctctaaag cccaacctt c 21

<210> 23

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 23

tgccgaagag gttcagtgc

19

<210> 24

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 24

gccacagtgg tactgtccag at

22

<210> 25

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 25

gctgcaagtt ctccacattg a

21

<210> 26

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 26

cagccgcagg tgaaacac

18

<210> 27

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 27

tggctttgaa ctcagggtca

20

<210> 28

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 28

cggatgcacc tcgtagacag

20

<210> 29

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 29

cggcaacctg gtagtgagtg

20

<210> 30

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 30

cgcagctcct tgtaaacttc ag

22

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 31
cgggaaccta ccagcctatg 20

<210> 32
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 32
caggcaacag ggagtcattgt 20

<210> 33
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 33
tgggcatctg ggtgtcaa 18

<210> 34
<211> 19
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 34
cggctgcgat gaggaagta 19

<210> 35
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 35
gcccatctcc tgcttcttta gt 22

<210> 36

WO 00/23111

PCT/US99/24331

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 36

cgtggagatg gctctgatgt a

21

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24331

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1, 141.1, 155.1, 183.1; 435/6, 7.1, 7.23, 7.9, 91.2; 436/501, 504, 505, 547; 514/44; 536/23.5

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Medline, Biosis, Embase, Cancerlit, Scisearch, WPIDS, USPATFULL
search terms: CSG, cancer specific gene, cancer, diagnosis

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Database SCISEARCH, Accession Number 307617, OLSSON et al. Reverse transcriptase-polymerase chain reaction assays for prostate cancer. Urologic Clinics of North America. May 1997, Vol. 24 No. 2, pages 367-&.	1-6
Y	CHO-CHUNG et al. Antisense Oligonucleotides for the treatment of cancer. Current Opinion in Therapeutic Patents. 1993, Vol. 3, No. 12, pages 1737-1750, see entire document.	1-6
A,E	BUSSEMAKERS et al. DD3: A new prostate-specific gene, highly overexpressed in prostate cancer. Cancer Research. 01 December 1999, Vol. 59, No. 23, pages 5975-5979.	1-7

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

10 FEBRUARY 2000

Date of mailing of the international search report

07 MAR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GEETHA P. BANSAL

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/24331

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (7):

A61K 39/395, 48/00; C12P 19/34; C12Q 1/68; G01N 33/53, 33/574, 33/546, 33/567

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/130.1, 141.1, 155.1, 183.1; 435/6, 7.1, 7.23, 7.9, 91.2; 436/501, 504, 505, 547; 514/44; 536/23.5